

ENVIRONMENTAL IMPACT ASSESSMENT FOR THE PROPOSED HOTAZEL 2 SOLAR PV
FACILITY AND ASSOCIATED INFRASTRUCTURE, HOTAZEL, NORTHERN CAPE:

FAUNA & FLORA SPECIALIST EIA REPORT



Cape EAPrac



PRODUCED FOR CAPE EAPRAC

BY



Simon.Todd@3foxes.co.za

February 2021

EXECUTIVE SUMMARY

Hotazel Solar Facility 2 (Pty) Ltd is proposing the establishment of a 100 MW commercial photovoltaic (PV) solar energy facility, called Hotazel 2, on the Remaining Extent (Portion 0) of the farm York A 279, situated in the District of Hotazel in the Northern Cape Province. The development is currently in the EIA Phase and 3Foxes Biodiversity Solutions has been appointed to provide a specialist terrestrial biodiversity EIA study of the development site as part of the authorization process.

A full field assessment as well as a desktop review of the available ecological information for the area was conducted in order to identify and characterise the ecological features of the site. The site falls within the Kathu Bushveld vegetation type, which is a relatively localised vegetation type for an arid area, but has not been significantly impacted by transformation and is classified as Least Threatened. The site has a relatively high abundance of *Vachellia erioloba* and *Vachellia haematoxylon*, which are nationally protected tree species. *Vachellia haematoxylon* is particularly common and approximately 3000-6000 individuals would potentially be lost as a result of the development. The extent of habitat loss (ca. 270 ha) associated with the development is however not seen as being highly significant for these species and as such no additional specific mitigation in this regard is considered necessary.

Cumulative impacts associated with the development are a concern given the development pressure in the wider Hotazel-Kathu area. The loss of 270ha of habitat associated with the development is however not considered highly significant given the spatial context of the site adjacent to mining, railway and road footprint areas. In terms of potential losses to landscape connectivity, the location of the site in an impacted area indicates that it is not likely an area that is important for faunal movement. As such, the overall cumulative impact of the development is considered to be low. This is supported by the fact that the area does not fall within a CBA or within a national or provincial protected area expansion strategy focus area.

Impact Statement

The development footprint of Hotazel 2 is restricted largely to low and moderately sensitive habitat typical of the wider Hotazel area. The affected area is considered suitable for development and there are no impacts associated with Hotazel 2 that cannot be mitigated to a low level. As such there are no fatal flaws or high post-mitigation impacts that should prevent the development from proceeding. Based on the layout provided for the assessment, Hotazel 2 can be supported from a terrestrial ecology point of view.

The Hotazel 2 Solar Grid Connection with associated infrastructure is likely to generate low impacts on fauna and flora after mitigation. Although all three grid connection alternatives are considered acceptable, Option 2 would be most preferred followed by Option 3 and then Option 1. Option 1 is preferred by the developer and while it would generate higher impacts than the other two options, the residual impact of this option would be low. No high impacts

that cannot be avoided were observed and from a flora and terrestrial fauna perspective, there are no reasons to oppose the development of the grid connection and associated infrastructure.

CONTENTS

Executive Summary2

Contents4

Short CV/Summary of Expertise – Simon Todd6

Specialist Declaration7

1 INTRODUCTION8

Scope of Study8

1.1 Assessment Approach & Philosophy 10

1.2 Relevant Aspects of the Proposed Development 12

2 METHODOLOGY 14

2.1 Data Sourcing and Review 14

2.2 Sensitivity Mapping & Assessment..... 15

2.3 Site Visit and Field Assessment..... 15

2.4 Sampling Limitations and Assumptions 16

3 DESCRIPTION OF THE AFFECTED ENVIRONMENT- BASELINE 17

3.1 Broad-Scale Vegetation Patterns 17

3.2 Habitats & Plant Communities..... 18

3.3 Listed and Protected Plant Species 21

3.4 Faunal Communities 22

 3.4.1 Mammals 22

 3.4.2 Reptiles 23

 3.4.3 Amphibians 24

3.5 Critical Biodiversity Areas & Broad-Scale Processes 25

3.6 Current Baseline & Cumulative Impact..... 26

3.7 Site Sensitivity Assessment 27

4 IDENTIFICATION & NATURE OF IMPACTS..... 28

4.1 Identification of Impacts to be Assessed 28

5 ASSESSMENT OF IMPACTS..... 29

5.1 Hotazel 2 PV Development 30

 5.1.1 Planning & Construction Phase..... 30

 5.1.2 Operational Phase Impacts..... 31

 5.1.3 Decommissioning Phase Impacts 32

 5.1.4 Cumulative Impacts 34

5.2 Hotazel 2 Grid Connection 37

5.2.1	Planning & Construction Phase.....	37
5.2.2	Operational Phase.....	39
5.2.3	Decommissioning Phase.....	40
6	CONCLUSION & RECOMMENDATIONS.....	42
7	REFERENCES	43
8	Annex 1. List of Plant Species	44
9	Annex 2. List of Mammals	48
10	Annex 2. List of Reptiles	50
11	Annex 3. List of Amphibians	52
12	Annex 3. Results of DEA Screening Tool.....	53

SHORT CV/SUMMARY OF EXPERTISE – SIMON TODD



Simon Todd is Director and principal scientist at 3Foxes Biodiversity Solutions and has over 20 years of experience in biodiversity measurement, management and assessment. He has provided specialist ecological input on more than 200 different developments distributed widely across the country. This includes input on the Wind and Solar SEA (REDZ) as well as the Eskom Grid Infrastructure (EGI) SEA and Karoo Shale Gas SEA. He is on the National Vegetation Map Committee as representative of the Nama and Succulent Karoo Biomes. Simon Todd is a recognised ecological expert and is a past chairman and current deputy chair of the Arid-Zone Ecology Forum. He is registered with the South African Council for Natural Scientific Professions (No. 400425/11).

A selection of recent work is as follows:

Strategic Environmental Assessments

- Co-Author. Chapter 7 - Biodiversity & Ecosystems - Shale Gas SEA. CSIR 2016.
- Co-Author. Chapter 1 Scenarios and Activities – Shale Gas SEA. CSIR 2016.
- Co-Author – Ecological Chapter – Wind and Solar SEA. CSIR 2014.
- Co-Author – Ecological Chapter – Eskom Grid Infrastructure SEA. CSIR 2015.
- Contributor – Ecological & Conservation components to SKA SEA. CSIR 2017.


Recent Specialist Ecological Studies in the Vicinity of the Current Site

- Kathu Solar PV Facility. Fauna and Flora EIA Process. Cape EAPrac 2015.
- Mogobe Solar PV Facility. Fauna and Flora EIA Proces. Cape EAPrac 2015.
- Logoko Solar PV Facility. Fauna and Flora EIA Proces. Cape EAPrac 2015.
- RE Capital 10 Solar Power Plant, Postmasburg. Fauna and Flora EIA Proces. Cape EAPrac 2015.
- Walk-through study of Kumba Iron Ore expansion area at Dingleton, Northern Cape. MSA Group. 2017.
- Adams PV Project – EIA process and follow-up vegetation survey. Aurora Power Solutions. 2016.
- Mamatwane Compilation Yard. Fauna and Flora EIA process. ERM. 2013.
- Olifantshoek-Emil 132kV power line. Fauna and Flora BA process. Savannah Environmental 2017.

SPECIALIST DECLARATION

I, ..Simon Todd....., as the appointed independent specialist, in terms of the 2014 EIA Regulations, hereby declare that I:

-
- I act as the independent specialist in this application;
- I perform the work relating to the application in an objective manner, even if this results in views and findings that are not favourable to the applicant;
- regard the information contained in this report as it relates to my specialist input/study to be true and correct, and do not have and will not have any financial interest in the undertaking of the activity, other than remuneration for work performed in terms of the NEMA, the Environmental Impact Assessment Regulations, 2014 as amended and any specific environmental management Act;
- I declare that there are no circumstances that may compromise my objectivity in performing such work;
- I have expertise in conducting the specialist report relevant to this application, including knowledge of the Act, Regulations and any guidelines that have relevance to the proposed activity;
- I will comply with the Act, Regulations and all other applicable legislation;
- I have no, and will not engage in, conflicting interests in the undertaking of the activity;
- I have no vested interest in the proposed activity proceeding;
- I undertake to disclose to the applicant and the competent authority all material information in my possession that reasonably has or may have the potential of influencing - any decision to be taken with respect to the application by the competent authority; and - the objectivity of any report, plan or document to be prepared by myself for submission to the competent authority;
- I have ensured that information containing all relevant facts in respect of the specialist input/study was distributed or made available to interested and affected parties and the public and that participation by interested and affected parties was facilitated in such a manner that all interested and affected parties were provided with a reasonable opportunity to participate and to provide comments on the specialist input/study;
- I have ensured that the comments of all interested and affected parties on the specialist input/study were considered, recorded and submitted to the competent authority in respect of the application;
- all the particulars furnished by me in this specialist input/study are true and correct; and
- I realise that a false declaration is an offence in terms of regulation 48 and is punishable in terms of section 24F of the Act.

Signature of the specialist:  _____

Name of Specialist: ____ Simon Todd _____

Date: ____ 20 February 2021 _____

1 INTRODUCTION

Hotazel Solar Facility 2 (Pty) Ltd is proposing the establishment of a commercial photovoltaic (PV) solar energy facility, called Hotazel 2, on the Remaining Extent (Portion 0) of the farm York A 279, situated in the District of Hotazel in the Northern Cape Province. Hotazel 2 is to consist of solar PV technology with fixed, single or double axis tracking mounting structures, with a net generation (contracted) capacity of 100 MW_{AC} (MegaWatts), as well as associated infrastructure, with an estimated maximum footprint of ± 270 ha. Hotazel Solar Facility 2 (Pty) Ltd has appointed Cape EAPrac to function as the Environmental Assessment Practitioner (EAP) and to undertake the required application for environmental authorisation for the above development. The development is currently in the EIA Phase and 3Foxes Biodiversity Solutions has been appointed to provide a specialist fauna and flora study of the development site as part of the EIA process.

The purpose of the Hotazel 2 Terrestrial Biodiversity EIA Phase Report is to describe and detail the ecological features of the proposed PV project site, provide an assessment of the ecological sensitivity of the site, and identify the likely impacts associated with the development of the site as a solar PV facility. A full field assessment as well as a desktop review of the available ecological information for the area were conducted in order to identify and characterise the ecological features of the site. This information is used to derive an ecological sensitivity map which has been used to inform the layout of the development. An assessment of likely impacts associated with the construction, operation, and decommissioning phases of the development is provided. A variety of avoidance and mitigation measures associated with each identified impact are recommended to reduce the likely impact of the development, which should be included in the EMP for the development. The full scope of study is detailed below.

SCOPE OF STUDY

The scope of the study includes the following activities

- a description of the environment that may be affected by the activity and the manner in which the environment may be affected by the proposed project
- a description and evaluation of environmental issues and potential impacts (incl. using direct, indirect and cumulative impacts) that have been identified
- a statement regarding the potential significance of the identified issues based on the evaluation of the issues/impacts
- an indication of the methodology used in determining the significance of potential environmental impacts
- an assessment of the significance of direct indirect and cumulative impacts in terms of the following criteria:

- the nature of the impact, which shall include a description of what causes the effect, what will be affected, and how it will be affected
 - the extent of the impact, indicating whether the impact will be local (limited to the immediate area or site of development), regional, national or international
 - the duration of the impact, indicating whether the lifetime of the impact will be of a short-term duration (0-5 years), medium-term (5- 15 years), long-term (> 15 years, where the impact will cease after the operational life of the activity), or permanent
 - the probability of the impact, describing the likelihood of the impact actually occurring, indicated as improbable (low likelihood) probable (distinct possibility), highly probable (most likely), or definite (impact will occur regardless of any preventable measures)
 - the severity/beneficial scale indicating whether the impact will be very severe/beneficial (a permanent change which cannot be mitigated/permanent and significant benefit with no real alternative to achieving this benefit), severe/beneficial (long-term impact that could be mitigated/long-term benefit), moderately severe/beneficial (medium- to long-term impact that could be mitigated/ medium- to long-term benefit), slight, or have no effect
 - the significance which shall be determined through a synthesis of the characteristics described above and can be assessed as low medium or high
 - the status which will be described as either positive, negative or neutral
 - the degree to which the impact can be reversed
 - the degree to which the impact may cause irreplaceable loss of resources
 - the degree to which the impact can be mitigated
- a description and comparative assessment of all alternatives
 - recommendations regarding practical mitigation measures for potentially significant impacts, for inclusion in the Environmental Management Programme (EMPr)
 - an indication of the extent to which the issue could be addressed by the adoption of mitigation measures
 - a description of any assumptions uncertainties and gaps in knowledge
 - an environmental impact statement which contains:
 - a summary of the key findings of the environmental impact assessment;
 - an assessment of the positive and negative implications of the proposed activity;
 - a comparative assessment of the positive and negative implications of identified alternatives.

General Considerations:

- Disclose any gaps in information or assumptions made.

- Identify recommendations for mitigatory measures to minimise impacts.
- Outline additional management guidelines.
- Provide monitoring requirements, mitigation measures and recommendations in a table format as input into the Environmental Management Programme (EMPr) for faunal related issues.

A description of the potential impacts of the development and recommended mitigation measures are to be provided, which will be separated into the following project phases:

- Preconstruction
- Construction
- Operational Phase

1.1 ASSESSMENT APPROACH & PHILOSOPHY

This assessment is conducted according to the 2014 EIA Regulations (Government Notice Regulation 982) in terms of the National Environmental Management Act (Act 107 of 1998) as amended (NEMA), as well as the recently promulgated notice issued in terms of NEMA, "*National Environmental Management Act, 1998 (Act No. 107 Of 1998): Procedures to be followed for the assessment and minimum criteria for reporting of identified environmental themes in terms of section 24(5)(a) and (h) and 44 of the National Environmental Management Act, 1998, when applying for environmental authorisation [G 43110 – GN 320]*". The applicable site verification report as required, is included under Annex 5 of this report and the required *Protocol for the assessment and reporting of environmental impacts on terrestrial animal species* is provided in Annex 7.

In terms of NEMA, this assessment demonstrates how the proponent intends to comply with the principles contained in Section 2 of NEMA, which amongst other things, indicates that environmental management should:

- In order of priority aim to: avoid, minimise or remedy disturbance of ecosystems and loss of biodiversity;
- Avoid degradation of the environment;
- Avoid jeopardising ecosystem integrity;
- Pursue the best practicable environmental option by means of integrated environmental management;
- Protect the environment as the people's common heritage;
- Control and minimise environmental damage; and
- Pay specific attention to management and planning procedures pertaining to sensitive, vulnerable, highly dynamic or stressed ecosystems.

These principles serve as guidelines for all decision-making concerning matters that may affect the environment. As such, it is incumbent upon the proponent to show how the proposed activities would comply with these principles and thereby contribute towards the achievement of sustainable development as defined by the NEMA.

In order to adhere to the above principles and best-practice guidelines, the following approach forms the basis for the study approach and assessment philosophy:

The study will include data searches, desktop studies, site walkovers / field survey of the property and baseline data collection, describing:

- A description of the broad ecological characteristics of the site and its surrounds in terms of any mapped spatial components of ecological processes and/or patchiness, patch size, relative isolation of patches, connectivity, corridors, disturbance regimes, ecotones, buffering, viability, etc.

In terms of **pattern**, the following will be identified or described:

Community and ecosystem level

- The main vegetation type, its aerial extent and interaction with neighbouring types, soils or topography
- Threatened or vulnerable ecosystems (*cf. SA vegetation map/National Spatial Biodiversity Assessment, fine-scale systematic conservation plans, etc*)

Species level

- Red Data Book (RDB) species (giving location if possible using GPS)
- The viability of an estimated population size of the RDB species that are present (include the degree of confidence in prediction based on availability of information and specialist knowledge, i.e. High=70-100% confident, Medium 40-70% confident, Low 0-40% confident)
- The likelihood of other RDB species, or species of conservation concern, occurring in the vicinity (include degree of confidence)

Fauna

- Describe and assess the terrestrial fauna present in the area that will be affected by the proposed development.
- Conduct a faunal assessment that can be integrated into the ecological study.
- Describe the existing impacts of current land use as they affect the fauna.
- Clarify species of special concern (SSC) and that are known to be:
 - endemic to the region;
 - that are considered to be of conservational concern;
 - that are in commercial trade (CITES listed species); or

- are of cultural significance.
- Provide monitoring requirements as input into the Environmental Management Programme (EMPr) for faunal related issues.

Other pattern issues

- Any significant landscape features or rare or important vegetation associations such as seasonal wetlands, alluvium, seeps, quartz patches or salt marshes in the vicinity.
- The extent of alien plant cover of the site, and whether the infestation is the result of prior soil disturbance such as ploughing or quarrying (alien cover resulting from disturbance is generally more difficult to restore than infestation of undisturbed sites).
- The condition of the site in terms of current or previous land uses.

In terms of **process**, the following will be identified or described:

- The key ecological “drivers” of ecosystems on the site and in the vicinity, such as fire.
- Any mapped spatial component of an ecological process that may occur at the site or in its vicinity (i.e. *corridors* such as watercourses, upland-lowland gradients, migration routes, coastal linkages or inland-trending dunes, and *vegetation boundaries* such as edaphic interfaces, upland-lowland interfaces or biome boundaries).
- Any possible changes in key processes, e.g. increased fire frequency or drainage/artificial recharge of aquatic systems.
- Furthermore, any further studies that may be required during or after the EIA process will be outlined.
- All relevant legislation, permits and standards that would apply to the development will be identified.
- The opportunities and constraints for development will be described and shown graphically on an aerial photograph, satellite image or map delineated at an appropriate level of spatial accuracy.

1.2 RELEVANT ASPECTS OF THE PROPOSED DEVELOPMENT

The project is to be located on the Remaining Extent (Portion 0) of the farm York A 279, situated in the District of Hotazel in the Northern Cape Province. The total footprint would be about 270ha, with access from the R31 (Figure 1). Hotazel 2 is to consist of solar photovoltaic (PV) technology with fixed, single or double axis tracking mounting structures, with a net generation (contracted) capacity of 100 MW_{AC} (MegaWatts), as well as associated infrastructure, which will include:

- On-site substation / collector switching station;

- Auxiliary buildings (gate-house and security, control centre, office, warehouse, canteen & visitors centre, staff lockers etc.);
- Inverter-stations, transformers and internal electrical reticulation (underground cabling);
- Access and internal road network;
- Laydown area;
- There are three options proposed to connect Hotazel 2 to the Eskom Hotazel Substation:
 - Option 1 (Preferred): Overhead 132kV powerline from the Hotazel 2 on-site substation/ collector switching station to the Eskom Hotazel substation.
 - Option 2: Via a loop in loop out (LILO) into the Hotazel-Eldoret 132kV line.
 - Option 3: Overhead 132kV powerline from the Hotazel 2 on-site substation/ collector switching station to the Hotazel Solar collector switching station.
- Rainwater tanks; and
- Perimeter fencing and security infrastructure.




Figure 1. Layout of Hotazel 2, showing the location of the facility within the site as well as the grid connection corridor to the Eskom Hotazel Substation.

2 METHODOLOGY

2.1 DATA SOURCING AND REVIEW

Published literature and data sources consulted during this study include the following:

Vegetation:

- Vegetation types and their conservation status were extracted from the South African National Vegetation Map (Mucina & Rutherford 2006 and 2018 update) as well as the National List of Threatened Ecosystems (2018), where relevant.
- Information on plant species recorded for the broad area around the site was extracted from the SANBI POSA database hosted by SANBI. The species list was derived from a considerably larger area than the study site, however, it is necessary to consider this data to ensure a conservative approach as well as to counter the fact that the site itself or the immediate surrounding area has not been well sampled in the past.
- The IUCN conservation status of identified species was extracted from the database and is based on the Threatened Species Programme, Red List of South African Plants (2020).

Ecosystem

- Critical Biodiversity Areas (CBAs) were extracted from the Northern Cape Critical Biodiversity Areas Map (Oosthuysen & Holness 2016).
- Freshwater and wetland information was extracted from the National Wetland Map produced as part of the 2018 NBA.
- Important catchments and protected areas expansion areas were extracted from the Northern Cape Protected Areas Expansion Strategy 2016 (NC-PAES).

Fauna

- Lists of mammals, reptiles and amphibians which are likely to occur at the site were derived based on distribution records from the literature and ADU Virtual Museum spatial database (<http://vmus.adu.org.za/>).
- Literature consulted includes Branch (1988) and Alexander and Marais (2007) for reptiles, Du Preez and Carruthers (2009) for amphibians, Friedmann and Daly (2004) and Skinner and Chimimba (2005) for mammals.
- Apart from the literature sources, additional information on fauna was extracted from the ADU web portal <http://vmus.adu.org.za>
- The faunal species lists provided are based on species which are known to occur in the broad geographical area, as well as a preliminary assessment of the availability and quality of suitable habitat at the site.
- The conservation status of mammals is based on the IUCN Red List Categories (EWT/SANBI 2016), while reptiles are based on the South African Reptile Conservation

Assessment (Bates et al. 2013) and amphibians on Minter et al. (2004) as well as the IUCN (2020).

2.2 SENSITIVITY MAPPING & ASSESSMENT

An ecological sensitivity map of the site was produced by integrating the available ecological and biodiversity information available in the literature and various spatial databases with mapping based on the satellite imagery of the site as well as personal knowledge of the site. This includes delineating different habitat units identified on the satellite imagery and assigning likely sensitivity values to the units based on their ecological properties, conservation value and the potential presence of species of conservation concern. The ecological sensitivity of the different units identified in the mapping procedure was rated according to the following scale:

Sensitivity	Description
Low	Units with a low sensitivity where there is likely to be a low impact on ecological processes and terrestrial biodiversity. This category represents transformed or natural areas where the impact of development is likely to be local in nature and of low significance with standard mitigation measures.
Medium	Areas of natural or previously transformed land where the impacts are likely to be largely local and the risk of secondary impacts such as erosion low. Development within these areas can proceed with relatively little ecological impact provided that appropriate mitigation measures are taken.
High	Areas of natural or transformed land where a high impact is anticipated due to the high biodiversity value, sensitivity or important ecological role of the area. Development within these areas is undesirable and should only proceed with caution as it may not be possible to mitigate all impacts appropriately.
Very High/No-Go	Critical and unique habitats that serve as habitat for rare/endangered species or perform critical ecological roles. These areas are essentially no-go areas from a developmental perspective and should be avoided as much as possible.

2.3 SITE VISIT AND FIELD ASSESSMENT

The site visit was conducted over 3 full days from the 18th-20th of April 2018. During the site visit, the different biodiversity features, habitat, and landscape units present at the site were identified and mapped in the field. Specific features visible on the satellite imagery of the site were also marked for field inspection and were verified and assessed during the site visit. Walk-through-surveys were conducted, within representative areas across the different

habitat units identified, and all plant and animal species observed were recorded. Conditions at the time of the site visit were excellent for the field assessment as it followed good rains in the area and the vegetation was actively growing with the majority of species present being in flower. As a result, the timing of the site visit has not compromised the study in any way and there are no significant limitations with regards to the vegetation assessment. In terms of small mammals, a transect consisting of 50 Sherman traps was set up on the site and included most of the habitats present in the site. The traps were set up every evening before sunset and checked each morning before 8am. Three camera traps were also set up on site along roads and near to one of the watering points on the site, in positions where fauna are most likely to pass. Active searches for reptiles and amphibians were also conducted within habitats likely to harbour or be important for such species. Although the duration of the field assessment was relatively short, it provided a reliable baseline of the typical fauna present on the site. Rare species which occasionally move through the area, are less likely to be encountered, but the expected presence of such species is assessed based on available literature, experience in the area and the quality and nature of the habitat present on-site.

2.4 SAMPLING LIMITATIONS AND ASSUMPTIONS

The current study consisted of a detailed field assessment as well as a desktop study, which serves to significantly reduce the limitations and assumptions of the study. For the current assessment, the vegetation was in an excellent condition for sampling at the time of the field assessment. Therefore, there are few limitations with regards to the vegetation sampling and the timing of the site visit. The plant species lists obtained from the field assessment are therefore considered comprehensive and reliable.

In terms of fauna, a number of activities and steps have been taken to obtain a reliable indication of the faunal community in the area. Sherman trapping for small mammals was conducted at the site in order to better characterise the small mammal community and while the sampling period was short, this provides a reliable insight into the dominant species present. Camera trapping was conducted at the site over several days and nights. Although this was a short period, it nevertheless provides an insight into the common species present at the site. Apart from the active searches that were conducted for reptiles and amphibians during the current study, additional species presence is inferred based on results obtained from the previous studies the consultant has conducted in the area. However, many fauna are difficult to observe in the field and their potential presence at the site is evaluated based on the literature and available databases. Many remote areas have not been well-sampled in the past with the result that the species lists derived from the available spatial databases for the area do not always adequately reflect the actual fauna present at the site. This is acknowledged as a limitation of the study however it is substantially reduced by the previous experience in the area. In order to further reduce this limitation, and ensure a conservative approach, the species lists derived for the site from the literature were obtained from an area

significantly larger than the study site and are likely to include a much wider array of species than those that actually occur at the site. This is a cautious and conservative approach which takes the study limitations into account.

3 DESCRIPTION OF THE AFFECTED ENVIRONMENT- BASELINE

3.1 BROAD-SCALE VEGETATION PATTERNS

According to the national vegetation map (Mucina & Rutherford 2006, SANBI 2018), the site is restricted to the Kathu Bushveld vegetation type. This vegetation unit occupies an area of 7 443 km² and extends from around Kathu and Dibeng in the south, through Hotazel, and to the Botswana border between Van Zylsrus and McCarthysrus. In terms of soils, the vegetation type is associated with aeolian red sand and surface calcrete and deep sandy soils of the Hutton and Clovelly soil forms. The main land types are Ah and Ae with some Ag. The Kathu Bushveld vegetation type is still largely intact and less than 2% has been transformed by mining activity. Therefore, it has been classified as Least Threatened. It is however, poorly conserved and does not currently fall within any formal conservation areas. Although no endemic species are restricted to this vegetation type, a number of Kalahari endemics are known to occur in this vegetation type such as *Vachellia luederitzii* var *luederitzii*, *Antheophora argentea*, *Megaloprotachne albescens*, *Panicum kalaharensense* and *Neuradopsis bechuanensis*. It is more fully described as it occurs at the site in the next section.

Other vegetation types that occur in the immediate area include Kuruman Thornveld to the east and Gordonias Duneveld to the west, neither of which is of conservation concern nor occur within the site.




Figure 2. Broad-scale overview of the vegetation in and around the Hotazel 2 site. The vegetation map is an extract of the national vegetation map as produced by Mucina and Rutherford (2006/2018), and also includes wetlands delineated by the NFEPA assessment (Nel et al. 2011).

3.2 HABITATS & PLANT COMMUNITIES

The vegetation on the proposed property consists of Bushveld with a well-developed grass layer and a variable-density tree layer. A feature of the property, which is also clearly visible from the aerial imagery, is the presence of *Senegalia mellifera* subsp. *detinens* bush clumps. As this is a bush encroaching species, it is considered to represent a symptom of degradation and the aggregations of trees present are thus not considered sensitive. Apart from the *Senegalia mellifera* subsp. *detinens* bush clumps, *Vachellia erioloba* and *Vachellia*

haematoxylon are also dominant species across large parts of the property. The density of these species increases towards the western boundary, however, this is outside of the proposed footprint for Hotazel 2. The grass layer is fairly homogenous across the site and there is not a lot a variation in the grass layer which can be ascribed to the consistent sandy substrate. Apart from the above dominant trees, other common woody species present at the site include *Zizyphus mucronata*, *Gymnosporia buxifolia*, *Senegalia haematoxylon* subsp. *detinens*, *Searsia ciliata*, *Ehretia rigida* subsp. *rigida*, *Diospyros lycioides* subsp. *lycioides* and *Grewia flava*. The grass layer is dominated by *Schmidtia pappophoroides*, *Aristida meridionalis*, *Aristida stipitata* subsp. *stipitata*, *Stipagrostis uniplumis* var. *uniplumis*, *Stipagrostis obtusa*, *Cynodon dactylon*, *Enneapogon desvauxii*, *Eragrostis lehmanniana* and *Aristida congesta* subsp. *congesta*. The density and diversity is shrubs is fairly low but includes *Asparagus laricinus*, *Asparagus retrofractus*, *Felicia muricata* subsp. *cinerascens*, *Pentzia calcarea*, *Vachellia hebeclada*, *Hermannia tomentosa*, *Gnidia polycephala* and *Lantana rugosa*. Due to the good rains preceding the site visit, forbs were abundant and included *Dicoma schinzii*, *Geigeria ornativa*, *Elephantorrhiza elephantina*, *Indigofera daleoides* var. *daleoides* and *Gisekia pharnacioides* var. *pharnacioides*.



Figure 3. The south-eastern corner of the development area, showing the slightly higher density of trees in this area which are comprised mostly of *Senegalia mellifera* with occasional *Vachellia haematoxylon* and *V.erioloba*.



Figure 4. Typical vegetation of the site with scattered *Vachellia haematoxylon* and *V.erioloba* within a matrix of grasses.



Figure 5. The typical vegetation within the proposed Hotazel 2 footprint consists of a well-developed grass layer with occasional *Vachellia haematoxylon* and *Senegalia mellifera*. Some of these more open areas have likely been achieved through the application of herbicides to eliminate bush-thickening species such as *Senegalia mellifera*.

3.3 LISTED AND PROTECTED PLANT SPECIES

Two NFA-protected tree species occur at the site and within the Hotazel 2 footprint, *Vachellia erioloba* and *Vachellia haematoxylon*. The density of both species is fairly high across the whole site and it would not be possible to avoid impact on these species. Although *Vachellia erioloba* has a higher density in some parts of the site, *Vachellia haematoxylon* is widely distributed across the site and there are no areas where this species does not occur to some degree. The density of *Vachellia haematoxylon* at the site varies from less than 10 trees/ha to approximately 30 trees/ha in the higher density areas. The Hotazel 2 footprint is located within an area with below-average density of *Vachellia erioloba* and *Vachellia haematoxylon*. However, due to the consistent presence of these species across the site, a few thousand trees at minimum would likely be lost as a result of the development. Both *Vachellia erioloba* and *Vachellia haematoxylon* are however very common in the area and their loss from the development area would not compromise their local populations. Devils' Claw *Harpagophytum procumbens* is common at the site, especially in the west of the site. It is likely that several hundred individuals of this species would be affected as a result of the development of Hotazel 2, but as *H.procumbens* is common in the area, the local population would not be significantly affected by the development.



Figure 6. Devils' Claw is common at the site, but the density within the proposed Hotazel 2 footprint is relatively low and the local population would not be compromised by the loss of the affected individuals.

3.4 FAUNAL COMMUNITIES

3.4.1 Mammals

The mammalian community at the site is likely to be of moderate diversity; although more than 50 species of terrestrial mammals are known from the wider area, the extent and habitat diversity of the site is too low to support a very wide range of mammals. Species observed or otherwise confirmed present at the site (Figure 7, Figure 8) include Aardvark, Cape Porcupine, Springhare, South African Ground Squirrel, Scrub Hare, Vervet Monkey, Small-spotted Genet, Yellow Mongoose, Slender Mongoose, Black-Backed Jackal, Steenbok, Duiker and Kudu. Small mammals trapped in the area during the current or previous site visits include Desert Pygmy Mouse *Mus indutus*, Multimammate Mouse *Mastomys coucha*, Bushveld Gerbil *Tatera leucogaster*, Hairy footed Gerbil *Gerbillurus paebe*, Pouched Mouse *Saccostomus campestris* and Grey Climbing Mouse *Dendromus melanotis*.



Figure 7. Species observed with the camera traps at the site include from top left, Common Duiker, Steenbok, Black-backed Jackal and Scrub Hare.

Five red-listed terrestrial mammal species potentially occur in the area; these are the Brown Hyena *Hyaena brunnea* (Near Threatened), Black-footed Cat *Felis nigripes* (Vulnerable), Leopard *Panthera pardus* (VU), Ground Pangolin *Smutsia temminckii* (Vulnerable) and South African Hedgehog *Atelerix frontalis* (Vulnerable). The Leopard and Brown Hyaena are not likely to occur in the area on account of the agricultural land-use in the area which is not usually conducive to the persistence of large carnivores. The Black-footed Cat is a secretive species which occurs across most of the Northern Cape and as such is likely to be present in the broad area given that the habitat is seen as broadly suitable. The Hedgehog and Ground Pangolin may also occur in the area at typically low density. Given the extensive national ranges of these species, the impact of the development on habitat loss for these species would be minimal and a long-term impact on these species would be unlikely.



Figure 8. Small mammals observed at the site include from top left, Pouched Mouse, Hairy-footed Gerbil and Bushveld Gerbil.

3.4.2 Reptiles

The Hotazel 2 site lies in or near the distribution range of more than 50 reptile species, although many of these are unlikely to occur on site, as it is restricted to sandy substrates and does not include rocky habitat or other habitats that are important for reptiles (Appendix 3). No species of conservation concern are known to occur in the area. The habitat diversity

within the study area is relatively low. As a result, the number of reptile species present within the site is likely to be relatively low.

Species observed at the site or in the area in the past include Serrated Tent Tortoise *Psammobates oculifer* (Figure 9), Cape Cobra *Naja nivea*, Ground Agama *Agama aculeata*, Spotted Sand Lizard *Pedioplanis lineocellata*, Variable Skink *Trachylepis varia*, Bibron's Blind Snake *Afrotyphlops bibronii*, Western Rock Skink *Mabuya sulcata sulcata*, Cape Gecko *Lygodactylus capensis capensis*, Speckled Rock Skink *Trachylepis punctatissima*, Striped Skaapsteker *Psammophylax tritaeniatus* and Boomslang *Dispholidus typus typus*. Impacts on reptiles are likely to be restricted largely to habitat loss within the development footprint. This is likely to be of local significance only as there are no very rare species or specialised habitats present within the proposed footprint.



Figure 9. The Serrated Tent Tortoise *Psammobates oculifer* was observed at the site.

3.4.3 Amphibians

The site lies within or near the range of 10 amphibian species, indicating that the site potentially has a moderately diverse frog community for an arid area. There is no natural permanent water or artificial earth dams within the site that would represent suitable breeding habitat for most of these species. Given the paucity of permanent water at the site, only those species which are relatively independent of water are likely to occur in the area. Species observed in the area include Eastern Olive Toad *Amietophrynus garmani* and Bushveld Rain Frog *Breviceps adspersus*, both of which are likely to occur at the site. There is no standing water on the site that could be used by amphibians for breeding purposes.

The only species of conservation concern which occurs in the wider area is the Giant Bullfrog *Pyxicephalus adspersus*. The site lies at the margin of the known distribution of this species and it has not been recorded from any of the quarter degree squares around the site, suggesting that it is unlikely to occur at the site. Impacts on amphibians are however likely to be low and restricted largely to habitat loss during construction.

3.5 CRITICAL BIODIVERSITY AREAS & BROAD-SCALE PROCESSES

An extract of the Northern Cape Critical Biodiversity Areas map for the study area is depicted below in Figure 10. The site lies within an area classified as “Other natural areas” and is not classified as a CBA or ESA. There are no CBAs in close proximity to the site, indicating that the development does not pose a threat to any CBAs or other areas considered to be of significance from a broad-scale conservation planning perspective.




Figure 10. Extract of the Northern Cape Critical Biodiversity Areas map for the site and surrounds, showing that there are no CBAs in close proximity to the site.

3.6 CURRENT BASELINE & CUMULATIVE IMPACT

There are several other proposed PV facilities in the wider Hotazel area (Figure 11), this would include the approved Hotazel Solar facility within the same property as the current project. The only constructed project in the immediate vicinity is the 10MW Adams PV facility 15km south of the site. However, in the wider area, there are several constructed PV plants towards Kathu including the Kalahari Solar, Kathu Solar and Sishen Solar Farms. The total extent of the constructed plants in the wider area as far south as Kathu is approximately 1000ha. The already built solar power plants are considered to form part of the existing baseline for the area and represent existing impact. The 1000ha footprint of these is however small in comparison with the iron and manganese mines in the area, which, with an existing footprint of at least 12 000ha, are currently the major driver of habitat loss and transformation in the Kathu-Hotazel area. There are also several authorised developments in close vicinity to the Hotazel 2 site, raising the potential for cumulative impact in the area. However, the overall development pressure in the wider area is still low and the proximity of the current development to Hotazel, the road and railway line as well as existing mine footprint areas suggests that the site is not likely to be of high significance for landscape connectivity. Consequently, the overall extent of cumulative impact due to the solar energy development in the area is seen to be relatively low and the contribution of the current development to cumulative impact is seen as low and of local significance only. The specific contribution of the current development is up to 270ha.




Figure 11. Map of DEA registered renewable energy applications in the vicinity of the Hotazel

2 site.

3.7 SITE SENSITIVITY ASSESSMENT




Figure 12. Sensitivity map for study area, showing the location of the Hotazel 2 footprint and grid connection route.

The sensitivity map for the Hotazel study area is illustrated above in Figure 12. There is not a lot of variation in sensitivity across the site, with the main driver of differences being the density of protected trees such as *Vachellia erioloba* and *Vachellia haematoxylon*. The majority of the site is considered medium sensitivity. Apart from the protected trees, the study area has a low abundance of other species or features of conservation concern. The west of the site as well as a small area in the eastern corner of the site are considered

moderately high sensitivity on account of the slightly higher tree density in these areas. No no-go or very high sensitivity areas were observed at the site and while it is considered broadly suitable for development, the potential impact on protected tree species is a concern. The majority of the Hotazel 2 footprint is within an area with lower than average density of these protected trees. As a result, the negative impact on the local populations of these species would be relatively low. Although it is common practice to consider the number of individuals of protected trees impacted by a particular development, the ultimate concern should be around the extent of habitat loss resulting from the development within habitats and vegetation types which support these species. When considered in this light, the 270ha of habitat loss is not considered to represent a large amount of habitat loss for either *V.erioloba* or *V.haematoxylon* which are widely distributed and are the dominant species across large areas surrounding the study area.

In terms of the three grid connection options, Option 2, the loop in loop out (LILO) into the Hotazel-Eldoret 132kV line would generate the least impact as the required power lines would be very short (<100m). Option 3, which would be an overhead 132kV powerline from the Hotazel 2 on-site substation/collector switching station to the Hotazel Solar collector switching station, would also generate low impacts as it would connect to the adjacent substation within the site. These options are however contingent on other projects and Option 1, the preferred connection which would be an overhead 132kV powerline from the Hotazel 2 on-site substation/collector switching station to the Eskom Hotazel substation is independent of the other projects. This would however require an overhead line of approximately 6.7km and would generate the highest relative impact compared to the other alternatives. However, it would run adjacent to existing lines, and it would not generate any high impacts with the result that it is considered acceptable but not the most desirable grid connection alternative should the other alternatives be viable.

4 IDENTIFICATION & NATURE OF IMPACTS

In this section, the potential impacts and associated risk factors that may be generated by the development are identified.

4.1 IDENTIFICATION OF IMPACTS TO BE ASSESSED

In this section each of the potential impacts identified as being likely to occur as a result of the development is explored in context of the features and characteristics of the site and the likelihood that each impact would occur given the nature of the development.

Impacts on vegetation and protected plant species

Several protected species, which may be impacted by the development, occur at the site, most notably *Vachellia erioloba* and *V.haematoxylon*. Vegetation clearing during

construction will lead to the loss of currently intact habitat within the development footprint and is an inevitable consequence of the development. This impact will be assessed for the construction phase as this is when the impact will occur, although the consequences will persist for a long time after construction.

Direct faunal impacts

Increased levels of noise, pollution, disturbance and human presence during construction will be detrimental to fauna. Sensitive and shy fauna would move away from the area during the construction phase as a result of the noise and human activities present. Some slow-moving species would not be able to avoid the construction activities and might be killed. Some impact on fauna is highly likely to occur during construction as well as operation of the facility. Therefore, this impact will be assessed for both the construction and operational phase.

Reduced ability to meet conservation obligations & targets

The loss of unprotected vegetation types on a cumulative basis from the broad area may impact the country's ability to meet its conservation targets. Although the receiving vegetation type in the study area is classified as Least Threatened and is still more than 98% intact, it is a relatively restricted vegetation type for an arid area and is therefore vulnerable to cumulative impact. This impact is therefore assessed in light of the current development as well as any other developments in the surrounding area which would also contribute to cumulative impacts.

Impact on broad-scale ecological processes

Transformation of intact habitat on a cumulative basis would contribute to the fragmentation of the landscape and would potentially disrupt the connectivity of the landscape for fauna and flora and impair their ability to respond to environmental fluctuations. Due to the presence of a number of other renewable energy and mining developments in the area, this is a potential cumulative impact of the development that is assessed.

5 ASSESSMENT OF IMPACTS

The various identified impacts are assessed below for the different phases of the development. It is important to note that the assessment is based on the layouts as provided and any changes to the layout or project description could invalidate the assessment.

5.1 HOTAZEL 2 PV DEVELOPMENT

The following is an assessment of the Hotazel 2 facility, for the planning and construction and operational phase of the development.

5.1.1 Planning & Construction Phase

Impact 1. Impacts on vegetation and protected plant species resulting from construction activities

Nature of impact	Impacts on vegetation and listed or protected plant species resulting from construction activities							
	Spatial Extent	Duration	Intensity	Probability	Reversibility	Significance and Status		Confidence level
						Without Mitigation	With Mitigation	
Hotazel 2	Local	Long-Term	Medium	Definite	Low	Medium Negative	Medium Negative	High
Mitigation/Management Actions								
<ul style="list-style-type: none"> • Preconstruction walk-through of the facility in order to locate species of conservation concern that can be translocated (such as aloes) as well as comply with the Northern Cape Nature Conservation Act and DENC/DAFF permit conditions. • Vegetation clearing to commence only after walk through has been conducted and necessary permits obtained. • Preconstruction environmental induction for all construction staff on site to ensure that basic environmental principles are adhered to. This includes awareness of no littering, appropriate handling of pollution and chemical spills, avoiding fire hazards, minimizing wildlife interactions, remaining within demarcated construction areas etc. • Environmental Control Officer (EO) to provide supervision and oversight of vegetation clearing activities within sensitive areas. • Vegetation clearing to be kept to a minimum. No unnecessary vegetation to be cleared. • All construction vehicles should adhere to clearly defined and demarcated roads. No off-road driving to be allowed outside of the construction area. • Temporary lay-down areas should be located within previously transformed areas or areas that have been identified as being of low sensitivity. These areas should be rehabilitated after use. 								

Impact 2. Direct Faunal Impacts Due to Construction Activities

Nature of impact	Direct Faunal Impacts During Construction							
	Spatial Extent	Duration	Intensity	Probability	Reversibility	Significance and Status		Confidence level
						Without Mitigation	With Mitigation	
Hotazel 2	Local	Short- Term	Medium	High	High	Medium	Medium-Low Negative	High
Mitigation/Management Actions								
<ul style="list-style-type: none"> All personnel should undergo environmental induction with regards to fauna and, in particular, awareness about not harming or collecting species such as snakes, tortoises and owls, which are often persecuted out of superstition. Any fauna threatened by the construction activities should be removed to safety by the EO or appropriately qualified environmental officer. All construction vehicles should adhere to a low speed limit to avoid collisions with susceptible species such as snakes and tortoises. All hazardous materials should be stored in the appropriate manner to prevent contamination of the site. Any accidental chemical, fuel and oil spills that occur at the site should be cleaned up in the appropriate manner as related to the nature of the spill. If trenches need to be dug for water pipelines or electrical cabling, these should not be left open for extended periods of time as fauna may fall in and become trapped in them. Trenches which are standing open should have places where there are soil ramps allowing fauna to escape the trench. 								

5.1.2 Operational Phase Impacts

Impact 1. Faunal Impacts due to Operation

Nature of Impact	Faunal Impacts due to operational activities								
	Alternative	Spatial Extent	Duration	Intensity	Probability	Reversibility	Significance and Status		Confidence level
							Without Mitigation	With Mitigation	
Hotazel 2	Local	Long-term	Medium-Low	Moderate	High	Medium-Low Negative	Low-Negative	High	
Mitigation/Management Actions									
<ul style="list-style-type: none"> Any potentially dangerous fauna such as snakes or fauna threatened by the maintenance and operational activities should be removed to a safe location. If the site must be lit at night for security purposes, this should be done with downward-directed low-UV type lights (such as most LEDs), which do not attract insects. 									

- All hazardous materials should be stored in the appropriate manner to prevent contamination of the site. Any accidental chemical, fuel and oil spills that occur at the site should be cleaned up in the appropriate manner as related to the nature of the spill.
- All vehicles accessing the site should adhere to a low speed limit (30km/h max) to avoid collisions with susceptible species such as snakes and tortoises.
- If the facility is to be fenced, then no electrified strands should be placed within 30cm of the ground as some species such as tortoises are susceptible to electrocution from electric fences because they do not move away when electrocuted but rather adopt defensive behaviour and are killed by repeated shocks. Alternatively, the electrified strands should be placed on the inside of the fence and not the outside as is the case on the majority of already constructed PV plants.

5.1.3 Decommissioning Phase Impacts

Impact 1. Direct Faunal Impacts Due to Decommissioning Activities

Nature of impact	Direct Faunal Impacts During Decommissioning							
	Spatial Extent	Duration	Intensity	Probability	Reversibility	Significance and Status		Confidence level
						Without Mitigation	With Mitigation	
Hotazel 2	Local	Short-Term	Medium	High	High	Medium-Low Negative	Low	High
Mitigation/Management Actions								
<ul style="list-style-type: none"> • All personnel should undergo environmental induction with regards to fauna and, in particular, awareness about not harming or collecting species such as snakes, tortoises and owls, which are often persecuted out of superstition. • Any fauna threatened by the decommissioning activities should be removed to safety by the EO or appropriately qualified environmental officer. • Any vehicles on-site should adhere to a low speed limit to avoid collisions with susceptible species such as snakes and tortoises. • All hazardous materials should be stored in the appropriate manner to prevent contamination of the site. Any accidental chemical, fuel and oil spills that occur at the site should be cleaned up in the appropriate manner as related to the nature of the spill. • If trenches need to be dug to remove water pipelines or electrical cabling, these should not be left open for extended periods of time as fauna may fall in and become trapped in them. Trenches which are standing open should have places where there are soil ramps allowing fauna to escape the trench. 								

Impact 2. Habitat Degradation due to Erosion and Alien Plant Invasion

Nature of impact	Habitat Degradation due to Erosion and Alien Plant Invasion							
	Spatial Extent	Duration	Intensity	Probability	Reversibility	Significance and Status		Confidence level
						Without Mitigation	With Mitigation	
Hotazel 2	Local	Long-Term	Medium	Medium	Medium	Medium-Low Negative	Low	High
Mitigation/Management Actions								
<ul style="list-style-type: none"> Erosion management at the site should take place according to the Erosion Management Plan and Rehabilitation Plan. This should make provision for monitoring of the site for at least 5 years after decommissioning. All erosion problems observed should be rectified as soon as possible, using the appropriate erosion control structures and revegetation techniques. There should be follow-up rehabilitation and revegetated of any remaining bare areas with indigenous perennial shrubs, grasses and trees from the local area. Alien management at the site should take place according to the Alien Invasive Management Plan. This should make provision for alien monitoring and management for at least 5 years after decommissioning. Regular (annual) monitoring for alien plant during operation to ensure that no erosion problems have developed as result of the disturbance, as per the Alien Management Plan for the project. Woody aliens should be controlled on at least an annual basis using the appropriate alien control techniques as determined by the species present. 								

5.1.4 Cumulative Impacts

The following are the cumulative impacts that are assessed as being a likely consequence of the development of the Hotazel 2 Facility. These are assessed in context of the extent of the current site, other developments in the area as well as general habitat loss and transformation resulting from mining and other activities in the area.

Cumulative Impact 1. Reduced ability to meet conservation obligations & targets due to cumulative habitat loss

Nature of impact	Reduced ability to meet conservation obligations & targets due to cumulative habitat loss							
Alternative	Spatial Extent	Duration	Intensity	Probability	Reversibility	Significance and Status		Confidence level
						Without Mitigation	With Mitigation	
Hotazel 2	Regional	Long-Term	Low	Low	Moderate	Low Negative	Low Negative	Moderate-High
<p>Mitigation/Management Actions</p> <ul style="list-style-type: none"> • The development footprint should be kept to a minimum and natural vegetation should be encouraged to return to disturbed areas. • An open space management plan should be developed for the site, which should include management of biodiversity within the fenced area, as well as that in the adjacent rangeland. 								

Cumulative Impact 2. Impact on broad-scale ecological processes due to cumulative loss and fragmentation of habitat

Nature of Impact	Impact on broad-scale ecological processes due to cumulative loss and fragmentation of habitat							
Alternative	Spatial Extent	Duration	Intensity	Probability	Reversibility	Significance and Status		Confidence level
						Without Mitigation	With Mitigation	
Hotazel 2	Regional	Long-Term	Medium	Moderate	Low	Medium-Low Negative	Low Negative	Moderate-High
<p>Mitigation/Management Actions</p> <ul style="list-style-type: none"> Minimise the development footprint as far as possible. A cover of indigenous grasses should be encouraged and maintained within the facility. This prevents the invasion of weeds and is the easiest to manage in the long-term. Furthermore, if possible, the grasses can be maintained low through livestock (sheep) grazing which is being successfully used at some existing PV facilities (see Figure 13 below for an operational example). The facility should be fenced off in a manner which allows small fauna to pass through the facility. In practical terms this means that the facility should be fenced-off to include only the developed areas and should include as little undeveloped ground or natural veld as possible. In addition, there should not be electrified ground-strands present within 30cm of the ground and the electrified strands should be located on the inside of the fence and not the outside. Furthermore, the fence should be a single layer fence and not a double fence with a large gap between. Images of suitable fencing types from existing PV facilities are available on request. 								



Figure 13. Sheep grazing within different PV plants in the Northern Cape.

5.2 HOTAZEL 2 GRID CONNECTION

The following is an assessment of the Grid Connection for Hotazel 2, for the planning and construction and operational phases of the development.

5.2.1 Planning & Construction Phase

Impact 1. Impacts on vegetation and listed or protected plant species resulting from power line construction activities

Impact Nature	Impacts on vegetation and listed or protected plant species resulting from power line construction activities							
Nature of impact	Spatial Extent	Duration	Intensity	Probability	Reversibility	Significance and Status		Confidence level
						Without Mitigation	With Mitigation	
Grid Connection Alt 1	Local	Long-Term	Moderate	High	Moderate	Medium-Low Negative	Low Negative	High
Grid Connection Alt 2	Local	Long-Term	Moderate	Low	High	Low Negative	Very Low Negative	High
Grid Connection Alt 3	Local	Long-Term	Moderate	Moderate	Moderate	Low Negative	Low Negative	High
Mitigation/Management Actions								
<ul style="list-style-type: none"> • Preconstruction walk-through of the power line route in order to locate species of conservation concern that can be translocated as well as comply with the Northern Cape Nature Conservation Act and DENC/DAFF permit conditions. • Construction and vegetation clearing to commence only after walk through has been conducted and necessary permits obtained. • No large woody species should be unnecessarily cleared from the power line servitude. It may be necessary to remove some individuals from the directly beneath the power line due to safety and operational concerns, however, within the servitude the presence of large woody species does not increase the fire risk and there are no valid reasons to remove such trees. If these are too tall and cause safety problems, they can be cut to a lower height rather than removed and as growth rate in arid areas is slow. It would take many years before such trees would need to be trimmed again. Such trees can be trimmed to 1m height if necessary although this would almost certainly result in the mortality of large <i>Vachellia erioloba</i> individuals. DAFF has a guideline available for tree clearing and trimming within power line servitudes which should serve as a guide. • Preconstruction environmental induction for all construction staff to ensure that basic environmental principles are adhered to. This includes awareness as to no littering, appropriate handling of pollution and chemical spills, avoiding fire hazards, minimizing wildlife interactions, remaining within demarcated construction areas etc. 								

Impact Nature	Impacts on vegetation and listed or protected plant species resulting from power line construction activities							
Nature of impact	Spatial Extent	Duration	Intensity	Probability	Reversibility	Significance and Status		Confidence level
						Without Mitigation	With Mitigation	
<ul style="list-style-type: none"> Vegetation clearing along the power line corridor should only be conducted where necessary and should not be cleared using herbicides or with a bulldozer. Vegetation can be cleared manually with bush cutters to 0.5m height where necessary. Temporary lay-down areas should be located within previously transformed areas or areas that have been identified as being of low sensitivity. 								

Impact 2. Faunal Impacts due to power line construction activities.

Impact Nature	Direct Faunal Impacts During Construction							
Alternative	Spatial Extent	Duration	Intensity	Probability	Reversibility	Significance and Status		Confidence level
						Without Mitigation	With Mitigation	
Grid Connection Alt 1	Local	Short- Term	Medium-Low	High	High	Medium-Low Negative	Low Negative	High
Grid Connection Alt 2	Local	Short- Term	Low	Low	High	Low Negative	Very Low Negative	High
Grid Connection Alt 3	Local	Short- Term	Low	Medium	High	Low Negative	Low Negative	High

Mitigation/Management Actions

- All personnel should undergo environmental induction with regards to fauna and in particular awareness about not harming or collecting species such as snakes, tortoises and owls which are often persecuted out of superstition.
- Any fauna threatened by the construction activities should be removed to safety by the EO or appropriately qualified environmental officer.
- All construction vehicles should adhere to a low speed limit to avoid collisions with susceptible species such as snakes and tortoises.
- All hazardous materials should be stored in the appropriate manner to prevent contamination of the site. Any accidental chemical, fuel and oil spills that occur at the site should be cleaned up in the appropriate manner as related to the nature of the spill.
- If holes or trenches need to be dug, these should not be left open for extended periods of time as fauna may fall in and become trapped in them. Holes should only be dug when they are required and should be used and filled shortly thereafter.

5.2.2 Operational Phase

Impact 1. Faunal Impacts due to Operation of the Grid Connection

Nature of Impact	Faunal Impacts due to operational activities							
Alternative	Spatial Extent	Duration	Intensity	Probability	Reversibility	Significance and Status		Confidence level
						Without Mitigation	With Mitigation	
Grid Connection Alt 1	Local	Long- Term	Medium-Low	Medium	High	Medium-Low Negative	Low Negative	High
Grid Connection Alt 2	Local	Long- Term	Low	Low	High	Low Negative	Very Low Negative	High
Grid Connection Alt 3	Local	Long- Term	Low	Medium	High	Low Negative	Low Negative	High
<p>Mitigation/Management Actions</p> <ul style="list-style-type: none"> Any potentially dangerous fauna such as snakes or fauna threatened by the maintenance and operational activities along the power line should be removed to a safe location if they pose a threat to staff, otherwise they should be left alone and allowed to move off on their own. If any parts of the grid connection infrastructure must be lit at night for security purposes, this should be done with downward-directed low-UV type lights (such as most LEDs or HPS bulbs), which attract fewer insects. All hazardous materials should be stored in the appropriate manner to prevent contamination of the site. Any accidental chemical, fuel and oil spills that occur at the site should be cleaned up in the appropriate manner as related to the nature of the spill. 								

5.2.3 Decommissioning Phase

Impact 1. Direct Faunal Impacts Due to Decommissioning Activities

Nature of impact	Direct Faunal Impacts During Decommissioning							
	Spatial Extent	Duration	Intensity	Probability	Reversibility	Significance and Status		Confidence level
						Without Mitigation	With Mitigation	
Grid Connection Alt 1	Local	Short- Term	Low	Medium	High	Medium-Low Negative	Low Negative	High
Grid Connection Alt 2	Local	Short- Term	Low	Low	High	Low Negative	Very Low Negative	High
Grid Connection Alt 3	Local	Short- Term	Low	Medium	High	Low Negative	Low Negative	High
<p>Mitigation/Management Actions</p> <ul style="list-style-type: none"> • All personnel should undergo environmental induction with regards to fauna and, in particular, awareness about not harming or collecting species such as snakes, tortoises and owls, which are often persecuted out of superstition. • Any fauna threatened by the decommissioning activities should be removed to safety by the EO or appropriately qualified environmental officer. • Any vehicles on-site should adhere to a low speed limit to avoid collisions with susceptible species such as snakes and tortoises. • All hazardous materials should be stored in the appropriate manner to prevent contamination of the site. Any accidental chemical, fuel and oil spills that occur at the site should be cleaned up in the appropriate manner as related to the nature of the spill. • If trenches need to be dug to remove pylons or electrical cabling, these should not be left open for extended periods of time as fauna may fall in and become trapped in them. Trenches which are standing open should have places where there are soil ramps allowing fauna to escape the trench. 								

Impact 2. Habitat Degradation due to Erosion and Alien Plant Invasion

Nature of impact	Habitat Degradation due to Erosion and Alien Plant Invasion							
	Spatial Extent	Duration	Intensity	Probability	Reversibility	Significance and Status		Confidence level
						Without Mitigation	With Mitigation	
Grid Connection Alt 1	Local	Long- Term	Medium-Low	Medium	High	Medium-Low Negative	Low Negative	High
Grid Connection Alt 2	Local	Long- Term	Low	Low	High	Low Negative	Very Low Negative	High
Grid Connection Alt 3	Local	Long- Term	Low	Medium	High	Low Negative	Low Negative	High
<p>Mitigation/Management Actions</p> <ul style="list-style-type: none"> • Erosion management along the power line should take place according to the Erosion Management Plan and Rehabilitation Plan. • There should be follow-up rehabilitation and revegetated of any remaining bare areas with indigenous perennial shrubs, grasses and trees from the local area. • Alien management in disturbed areas should take place according to the Alien Invasive Management Plan. This should make provision for alien monitoring and management for at least 3 years after decommissioning. • Woody aliens should be controlled on at least an annual basis using the appropriate alien control techniques as determined by the species present. 								

6 CONCLUSION & RECOMMENDATIONS

The vegetation of the Hotazel 2 site consists of Kathu Bushveld with a relatively high abundance of *Vachellia erioloba* and *Vachellia haematoxylon*. Although relatively large numbers of *Vachellia haematoxylon* (3000-6000) would potentially be lost as a result of the development, the extent to habitat loss (270 ha) is not seen as being highly significant for this species and is of local relevance only. Therefore, it is not seen as sufficient to warrant an offset or other similar off-site mitigation measures.

Cumulative impacts in the area are a concern due firstly to the mining activity that characterises the area and secondly due to the proliferation of solar energy development in the wider Hotazel-Kathu area. In terms of habitat loss, the affected Kathu Bushveld vegetation type is still approximately 90% intact and while this is not a very extensive vegetation type, the loss of 270ha of habitat is not considered highly significant, especially given the spatial context of the site adjacent to mining, railway and road footprint areas. In terms of potential losses to landscape connectivity, the location of the site in an impacted area indicates that it is not likely to be important for faunal movement. As such, the overall cumulative impact of the development is considered likely to be low. This is also supported by the fact that the area has not been identified as being a CBA or NPAES Focus Area.

Although all three grid connection alternatives are considered acceptable, the on-site grid connection options consisting of a loop-in loop-out connection to the 132kV line that traverses the site are preferable to the 6.7km connection to the Eskom Hotazel substation. However, regardless of which alternatives is ultimately used, there are no impacts associated with the grid connection development that are considered to be of high significance and which cannot be mitigated to an acceptable level.

Impact Statement

The development footprint of Hotazel 2 is restricted largely to low and moderately sensitive habitat typical of the wider Hotazel area. The affected area is considered suitable for development and there are no impacts associated with the Hotazel 2 facility that cannot be mitigated to a low level. As such, there are no fatal flaws or high post-mitigation impacts that should prevent the development from proceeding. Based on the layout provided for the assessment, Hotazel 2 can be supported from a terrestrial ecology point of view.

The Hotazel 2 Grid Connection with associated infrastructure is likely to generate low impacts on fauna and flora after mitigation. No high impacts that cannot be avoided were observed. Therefore, from a flora and terrestrial fauna perspective, there are no reasons to oppose the development of the grid connections and associated infrastructure.

7 REFERENCES

- Alexander, G. & Marais, J. 2007. *A Guide to the Reptiles of Southern Africa*. Struik Nature, Cape Town.
- Bates, M.F., Branch, W.R., Bauer, A.M., Burger, M., Marais, J., Alexander, G.J. & de Villiers, M. S. 2014. Atlas and Red List of the Reptiles of South Africa, Lesotho and Swaziland. Strelitzia 32. SANBI, Pretoria.
- Branch W.R. 1998. *Field guide to snakes and other reptiles of southern Africa*. Struik, Cape Town.
- Du Preez, L. & Carruthers, V. 2009. *A Complete Guide to the Frogs of Southern Africa*. Struik Nature., Cape Town.
- EWT & SANBI, 2016. Red List of Mammals of South Africa, Lesotho and Swaziland. EWT, Johannesburg.
- Marais, J. 2004. *Complete Guide to the Snakes of Southern Africa*. Struik Nature, Cape Town.
- Nel, J.L., Murray, K.M., Maherry, A.M., Petersen, C.P., Roux, D.J., Driver, A., Hill, L., Van Deventer, H., Funke, N., Swartz, E.R., Smith-Adao, L.B., Mbona, N., Downsborough, L. and Nienaber, S. (2011). Technical Report for the National Freshwater Ecosystem Priority Areas project. WRC Report No. K5/1801.
- Minter LR, Burger M, Harrison JA, Braack HH, Bishop PJ & Kloepfer D (eds). 2004. Atlas and Red Data book of the frogs of South Africa, Lesotho and Swaziland. SI/MAB Series no. 9. Smithsonian Institution, Washington, D.C.
- Mucina L. & Rutherford M.C. (eds) 2006. *The Vegetation of South Africa, Lesotho and Swaziland*. Strelitzia 19. South African National Biodiversity Institute, Pretoria.
- Oosthuysen, E. & Holness, S. 2016. Northern Cape Critical Biodiversity Areas (CBA) Map. Northern Cape Department of Environment and Nature Conservation & Nelson Mandela Metropolitan University. Available at SANBI BGIS <http://bgis.sanbi.org/>.
- Skinner, J.D. & Chimimba, C.T. 2005. The mammals of the Southern African Subregion. Cambridge University Press, Cambridge.

8 ANNEX 1. LIST OF PLANT SPECIES

List of plant species confirmed present at the Hotazel site during the course of the field assessment.

Family	Species	IUCN Status
ACANTHACEAE	<i>Barleria rigida</i>	LC
ACANTHACEAE	<i>Justicia puberula</i>	LC
AIZOACEAE	<i>Plinthus sericeus</i>	LC
AMARANTHACEAE	<i>Gomphrena celosioides</i>	LC
AMARANTHACEAE	<i>Hermbstaedia odorata</i> var. <i>odorata</i>	LC
AMARANTHACEAE	<i>Pupalia lappacea</i> var. <i>lappacea</i>	LC
AMARYLLIDACEAE	<i>Boophone disticha</i>	LC
ANACARDIACEAE	<i>Searsia ciliata</i>	LC
APOCYNACEAE	<i>Raphionacme velutina</i>	LC
ASPARAGACEAE	<i>Asparagus laricinus</i>	LC
ASPARAGACEAE	<i>Asparagus retrofractus</i>	LC
ASPHODELIACEAE	<i>Bulbine narcissifolia</i>	LC
ASTERACEAE	<i>Chrysocoma ciliata</i>	LC
ASTERACEAE	<i>Dicoma schinzii</i>	LC
ASTERACEAE	<i>Felicia muricata</i> subsp. <i>cinerascens</i>	LC
ASTERACEAE	<i>Gazania krebsiana</i> subsp. <i>krebsiana</i>	LC
ASTERACEAE	<i>Geigeria ornativa</i>	LC
ASTERACEAE	<i>Helichrysum zeyheri</i>	LC
ASTERACEAE	<i>Hertia pallens</i>	LC
ASTERACEAE	<i>Nolletia ciliaris</i>	LC
ASTERACEAE	<i>Osteospermum muricatum</i>	LC
ASTERACEAE	<i>Pegolettia retrofracta</i>	LC
ASTERACEAE	<i>Pentzia calcarea</i>	LC
ASTERACEAE	<i>Pentzia sphaerocephala</i>	LC
ASTERACEAE	<i>Pteronia incana</i>	LC
ASTERACEAE	<i>Rosenia humilis</i>	LC
ASTERACEAE	<i>Senecio inaequidens</i>	LC
ASTERACEAE	<i>Tarchonanthus camphoratus</i>	LC
ASTERACEAE	<i>Verbesina encelioides</i>	LC
BORAGINACEAE	<i>Ehretia rigida</i> subsp. <i>rigida</i>	LC
BORAGINACEAE	<i>Heliotropium ciliatum</i>	LC
CAPPARACEAE	<i>Cleome rubella</i>	LC
CELASTRACEAE	<i>Gymnosporia buxifolia</i>	LC
COMMELINACEAE	<i>Commelina africana</i> var. <i>africana</i>	LC
CUCURBITACEAE	<i>Acanthosicyos naudinianus</i>	LC
CUCURBITACEAE	<i>Coccinia sessilifolia</i>	LC
CUCURBITACEAE	<i>Cucumis africanus</i>	LC

CYPERACEAE	<i>Cyperus margaritaceus var. margaritaceus</i>	LC
CYPERACEAE	<i>Kyllinga alba</i>	LC
EBENACEAE	<i>Diospyros lycioides subsp. lycioides</i>	LC
ERIOSPERMACEAE	<i>Eriospermum sp.</i>	LC
EUPHORBIACEAE	<i>Tragia dioica</i>	LC
FABACEAE	<i>Senegalia hebeclada</i>	LC
FABACEAE	<i>Vachellia erioloba</i>	LC
FABACEAE	<i>Vachellia haematoxylon</i>	LC
FABACEAE	<i>Senegalia karroo</i>	LC
FABACEAE	<i>Senegalia haematoxylon subsp. detinens</i>	LC
FABACEAE	<i>Cyamopsis serrata</i>	LC
FABACEAE	<i>Elephantorrhiza elephantina</i>	LC
FABACEAE	<i>Indigofera daleoides var. daleoides</i>	LC
FABACEAE	<i>Lessertia pauciflora var. pauciflora</i>	LC
FABACEAE	<i>Melolobium exudans</i>	LC
FABACEAE	<i>Melolobium macrocalyx var. macrocalyx</i>	LC
FABACEAE	<i>Senna italica subsp. arachoides</i>	LC
FABACEAE	<i>Tephrosia burchellii</i>	LC
FABACEAE	<i>Tephrosia longipes subsp. longipes var. longipes</i>	LC
GERANIACEAE	<i>Monsonia angustifolia</i>	LC
GISEKIACEAE	<i>Gisekia pharnacioides var. pharnacioides</i>	LC
HYACINTHACEAE	<i>Dipcadi viride</i>	LC
HYACINTHACEAE	<i>Ledebouria ovatifolia</i>	LC
IRIDACEAE	<i>Babiana bainesii</i>	LC
LAMIACEAE	<i>Acrotome inflata</i>	LC
LAMIACEAE	<i>Leucas capensis</i>	LC
MALVACEAE	<i>Corchorus pinnatipartitus</i>	LC
MALVACEAE	<i>Grewia flava</i>	LC
MALVACEAE	<i>Hermannia comosa</i>	LC
MALVACEAE	<i>Hermannia jacobefolia</i>	LC
MALVACEAE	<i>Hermannia linnaeoides</i>	LC
MALVACEAE	<i>Hermannia tomentosa</i>	LC
MALVACEAE	<i>Hibiscus marlothianus</i>	LC
MALVACEAE	<i>Hibiscus pusillus</i>	LC
MALVACEAE	<i>Pavonia burchellii</i>	LC
MOLLUGINACEAE	<i>Hypertelis salsoloides</i>	LC
MOLLUGINACEAE	<i>Limeum aethiopicum var. intermedium</i>	LC
MOLLUGINACEAE	<i>Limeum argute carinatum var argute carinatum</i>	LC
MOLLUGINACEAE	<i>Limeum fenestratum var. fenestratum</i>	LC
MOLLUGINACEAE	<i>Limeum sulcatum var sulcatum</i>	LC
MOLLUGINACEAE	<i>Mollugo cerviana</i>	LC
OROBANCHACEAE	<i>Striga bilabiata subsp. bilabiata</i>	LC
OXALIDACEAE	<i>Oxalis depressa</i>	LC
OXALIDACEAE	<i>Oxalis lawsonii</i>	LC

PEDALIACEAE	<i>Sesamum triphyllum</i>	LC
PHYLLANTHACEAE	<i>Phyllanthus maderaspatensis</i>	LC
POACEAE	<i>Aristida adscensionis</i>	LC
POACEAE	<i>Aristida congesta subsp. congesta</i>	LC
POACEAE	<i>Aristida meridionalis</i>	LC
POACEAE	<i>Aristida stipitata subsp. graciliflora</i>	LC
POACEAE	<i>Aristida stipitata subsp. stipitata</i>	LC
POACEAE	<i>Brachiaria marlothii</i>	LC
POACEAE	<i>Cenchrus ciliaris</i>	LC
POACEAE	<i>Cymbopogon popischilli</i>	LC
POACEAE	<i>Cynodon dactylon</i>	LC
POACEAE	<i>Enneapogon cenchroides</i>	LC
POACEAE	<i>Enneapogon desvauxii</i>	LC
POACEAE	<i>Eragrostis biflora</i>	LC
POACEAE	<i>Eragrostis lehmanniana var. chaunantha</i>	LC
POACEAE	<i>Eragrostis nindensis</i>	LC
POACEAE	<i>Eragrostis obtusa</i>	LC
POACEAE	<i>Fingerhuthia africana</i>	LC
POACEAE	<i>Melinis repens subsp. repens</i>	LC
POACEAE	<i>Oropetium capense</i>	LC
POACEAE	<i>Pogonarthria squarrosa</i>	LC
POACEAE	<i>Schmidtia pappophoroides</i>	LC
POACEAE	<i>Stipagrostis obtusa</i>	LC
POACEAE	<i>Stipagrostis uniplumis var. uniplumis</i>	LC
POACEAE	<i>Tragus berteronianus</i>	LC
POLYGALACEAE	<i>Polygala seminuda</i>	LC
PORTULACACEAE	<i>Portulaca kermesina</i>	LC
PORTULACACEAE	<i>Talinum arnotii</i>	LC
RANUNCULACEAE	<i>Clematis brachiata</i>	LC
RHAMNACEAE	<i>Ziziphus mucronata subsp. mucronata</i>	LC
RUBIACEAE	<i>Kohautia caespitosa subsp. brachyloba</i>	LC
SCROPHULARIACEAE	<i>Aptosimum albomarginatum</i>	LC
SCROPHULARIACEAE	<i>Aptosimum elongatum</i>	LC
SCROPHULARIACEAE	<i>Aptosimum lineare var. lineare</i>	LC
SCROPHULARIACEAE	<i>Chaenostoma halimifolium</i>	LC
SCROPHULARIACEAE	<i>Jamesbrittenia atropurpurea subsp. atropurpurea</i>	LC
SCROPHULARIACEAE	<i>Peliostomum leuchorhizum</i>	LC
SCROPHULARIACEAE	<i>Selago mixta</i>	LC
SCROPHULARIACEAE	<i>Sutera griquensis</i>	LC
SOLANACEAE	<i>Datura stramonium</i>	LC
SOLANACEAE	<i>Lycium hirsutum</i>	LC
THYMELAEACEAE	<i>Gnidia polycephala</i>	LC
VAHLIACEAE	<i>Vahlia capensis subsp. vulgaris var. vulgaris</i>	LC
VERBENACEAE	<i>Chascanum pinnatifidum var. pinnatifidum</i>	LC

VERBENACEAE	<i>Lantana rugosa</i>	LC
ZYGOPHYLLACEAE	<i>Tribulus terrestris</i>	LC

9 ANNEX 2. LIST OF MAMMALS

List of mammals which have been observed or which are likely to occur in the vicinity of the Hotazel 2 site. Conservation status is from 2016 EWT/SANBI Red List.

Family	Scientific name	Common name	Red list category	Number of records
<i>Bathyergidae</i>	<i>Bathyergus janetta</i>	Namaqua Dune Mole-rat	Least Concern	1
<i>Bathyergidae</i>	<i>Cryptomys hottentotus</i>	Southern African Mole-rat	Least Concern	6
<i>Bathyergidae</i>	<i>Fukomys damarensis</i>	Damara Mole-rat	Least Concern	12
<i>Bovidae</i>	<i>Antidorcas marsupialis</i>	Springbok	Least Concern	7
<i>Bovidae</i>	<i>Oreotragus oreotragus</i>	Klipspringer	Least Concern	6
<i>Bovidae</i>	<i>Oryx gazella</i>	Gemsbok	Least Concern	16
<i>Bovidae</i>	<i>Raphicerus campestris</i>	Steenbok	Least Concern	9
<i>Bovidae</i>	<i>Sylvicapra grimmia</i>	Bush Duiker	Least Concern	8
<i>Bovidae</i>	<i>Tragelaphus strepsiceros</i>	Greater Kudu	Least Concern	12
<i>Canidae</i>	<i>Canis mesomelas</i>	Black-backed Jackal	Least Concern	10
<i>Canidae</i>	<i>Otocyon megalotis</i>	Bat-eared Fox	Least Concern	5
<i>Canidae</i>	<i>Vulpes chama</i>	Cape Fox	Least Concern	7
<i>Cercopithecidae</i>	<i>Papio ursinus</i>	Chacma Baboon	Least Concern	8
<i>Erinaceidae</i>	<i>Atelerix frontalis</i>	Southern African Hedgehog	Near Threatened	9
<i>Felidae</i>	<i>Caracal caracal</i>	Caracal	Least Concern	1
<i>Felidae</i>	<i>Felis nigripes</i>	Black-footed Cat	Vulnerable	3
<i>Felidae</i>	<i>Felis silvestris</i>	Wildcat	Least Concern	1
<i>Felidae</i>	<i>Panthera pardus</i>	Leopard	Vulnerable	4
<i>Gliridae</i>	<i>Graphiurus platyops</i>	Flat-headed African Dormouse	Data deficient	1
<i>Herpestidae</i>	<i>Cynictis penicillata</i>	Yellow Mongoose	Least Concern	2
<i>Herpestidae</i>	<i>Herpestes sanguineus</i>	Slender Mongoose	Least Concern	2
<i>Herpestidae</i>	<i>Suricata suricatta</i>	Meerkat	Least Concern	3
<i>Hyaenidae</i>	<i>Hyaena brunnea</i>	Brown Hyena	Near Threatened	12
<i>Hyaenidae</i>	<i>Proteles cristata</i>	Aardwolf	Least Concern	6
<i>Hystricidae</i>	<i>Hystrix africaeaustralis</i>	Cape Porcupine	Least Concern	16
<i>Leporidae</i>	<i>Lepus capensis</i>	Cape Hare	Least Concern	18
<i>Leporidae</i>	<i>Lepus saxatilis</i>	Scrub Hare	Least Concern	16

Fauna & Flora Specialist EIA Report

<i>Leporidae</i>	<i>Pronolagus rupestris</i>	Smith's Red Rock Hare	Least Concern	14
<i>Macroscelididae</i>	<i>Elephantulus intufi</i>	Bushveld Elephant Shrew	Least Concern	1
<i>Macroscelididae</i>	<i>Elephantulus myurus</i>	Eastern Rock Elephant Shrew	Least Concern	29
<i>Macroscelididae</i>	<i>Elephantulus rupestris</i>	Western Rock Elephant Shrew	Least Concern	37
<i>Macroscelididae</i>	<i>Macroscelides proboscideus</i>	Short-eared Elephant Shrew	Least Concern	1
<i>Manidae</i>	<i>Smutsia temminckii</i>	Ground Pangolin	Vulnerable	23
<i>Muridae</i>	<i>Aethomys chrysophilus</i>	Red Veld Aethomys	Least Concern	3
<i>Muridae</i>	<i>Aethomys namaquensis</i>	Namaqua Rock Mouse	Least Concern	171
<i>Muridae</i>	<i>Desmodillus auricularis</i>	Cape Short-tailed Gerbil	Least Concern	38
<i>Muridae</i>	<i>Gerbilliscus brantsii</i>	Highveld Gerbil	Least Concern	4
<i>Muridae</i>	<i>Gerbilliscus leucogaster</i>	Bushveld Gerbil	Least Concern	103
<i>Muridae</i>	<i>Gerbilliscus paeba</i>	Paeba Hairy-footed Gerbil	Least Concern	2
<i>Muridae</i>	<i>Gerbilliscus vallinus</i>	Brush-tailed Hairy-footed Gerbil	Least Concern	4
<i>Muridae</i>	<i>Mastomys coucha</i>	Southern African Mastomys	Least Concern	56
<i>Muridae</i>	<i>Mus (Nannomys) minutoides</i>	Southern African Pygmy Mouse	Least Concern	27
<i>Muridae</i>	<i>Otomys auratus</i>	Southern African Vlei Rat	Near Threatened	3
<i>Muridae</i>	<i>Parotomys brantsii</i>	Brants's Whistling Rat	Least Concern	1
<i>Muridae</i>	<i>Rhabdomys pumilio</i>	Xeric Four-striped Grass Rat	Least Concern	41
<i>Mustelidae</i>	<i>Ictonyx striatus</i>	Striped Polecat	Least Concern	2
<i>Mustelidae</i>	<i>Mellivora capensis</i>	Honey Badger	Least Concern	4
<i>Nesomyidae</i>	<i>Saccostomus campestris</i>	Southern African Pouched Mouse	Least Concern	45
<i>Orycteropodidae</i>	<i>Orycteropus afer</i>	Aardvark	Least Concern	4
<i>Pedetidae</i>	<i>Pedetes capensis</i>	South African Spring Hare	Least Concern	23
<i>Procaviidae</i>	<i>Procavia capensis</i>	Cape Rock Hyrax	Least Concern	15
<i>Sciuridae</i>	<i>Xerus inauris</i>	South African Ground Squirrel	Least Concern	16
<i>Soricidae</i>	<i>Crocidura cyanea</i>	Reddish-gray Musk Shrew	Least Concern	3
<i>Soricidae</i>	<i>Crocidura hirta</i>	Lesser Red Musk Shrew	Least Concern	12
<i>Suidae</i>	<i>Phacochoerus africanus</i>	Common Warthog	Least Concern	11

10 ANNEX 2. LIST OF REPTILES

List of reptiles which are likely to occur at the proposed Hotazel 2 site, based on the ReptileMap database. Conservation status is from Bates et al. (2014).

Family	Scientific name	Common name	Red list category	Number of records
Agamidae	<i>Agama aculeata aculeata</i>	Common Ground Agama	Least Concern	41
Agamidae	<i>Agama atra</i>	Southern Rock Agama	Least Concern	17
Amphisbaenidae	<i>Monopeltis mauricei</i>	Maurice's Worm Lizard	Least Concern	1
Amphisbaenidae	<i>Zygaspis quadrifrons</i>	Kalahari Dwarf Worm Lizard	Least Concern	4
Chamaeleonidae	<i>Chamaeleo dilepis dilepis</i>	Common Flap-neck Chameleon	Least Concern	8
Colubridae	<i>Dasypeltis scabra</i>	Rhombic Egg-eater	Least Concern	2
Colubridae	<i>Dispholidus typus typus</i>	Boomslang	Least Concern	3
Colubridae	<i>Philothamnus semivariiegatus</i>	Spotted Bush Snake	Least Concern	1
Colubridae	<i>Telescopus semiannulatus semiannulatus</i>	Eastern Tiger Snake	Least Concern	9
Cordylidae	<i>Karusasaurus polyzonus</i>	Karoo Girdled Lizard	Least Concern	7
Elapidae	<i>Aspidelaps scutatus scutatus</i>	Speckled Shield Cobra	Least Concern	4
Elapidae	<i>Dendroaspis polylepsis</i>	Black Mamba	Least Concern	1
Elapidae	<i>Naja nigricincta woodi</i>	Black Spitting Cobra	Least Concern	2
Elapidae	<i>Naja nivea</i>	Cape Cobra	Least Concern	4
Gekkonidae	<i>Chondrodactylus angulifer</i>	Giant Ground Gecko	Least Concern	4
Gekkonidae	<i>Chondrodactylus angulifer angulifer</i>	Common Giant Ground Gecko	Least Concern	9
Gekkonidae	<i>Chondrodactylus bibronii</i>	Bibron's Gecko	Least Concern	3
Gekkonidae	<i>Lygodactylus bradfieldi</i>	Bradfield's Dwarf Gecko	Least Concern	1
Gekkonidae	<i>Lygodactylus capensis capensis</i>	Common Dwarf Gecko	Least Concern	8
Gekkonidae	<i>Pachydactylus capensis</i>	Cape Gecko	Least Concern	14
Gekkonidae	<i>Pachydactylus rugosus</i>	Common Rough Gecko	Least Concern	1
Gekkonidae	<i>Pachydactylus wahlbergii wahlbergii</i>	Kalahari Ground Gecko	Least Concern	12
Gekkonidae	<i>Ptenopus garrulus garrulus</i>	Common Barking Gecko	Least Concern	12
Gerrhosauridae	<i>Gerrhosaurus flavigularis</i>	Yellow-throated Plated Lizard	Least Concern	1
Lacertidae	<i>Heliobolus lugubris</i>	Bushveld Lizard	Least Concern	23
Lacertidae	<i>Meroles squamulosus</i>	Common Rough-scaled Lizard	Least Concern	3
Lacertidae	<i>Nucras intertexta</i>	Spotted Sandveld Lizard	Least Concern	14

Fauna & Flora Specialist EIA Report

<i>Lacertidae</i>	<i>Pedioplanis lineocellata lineocellata</i>	Spotted Sand Lizard	Least Concern	37
<i>Lacertidae</i>	<i>Pedioplanis namaquensis</i>	Namaqua Sand Lizard	Least Concern	4
<i>Lamprophiidae</i>	<i>Aparallactus capensis</i>	Black-headed Centipede-eater	Least Concern	1
<i>Lamprophiidae</i>	<i>Atractaspis bibronii</i>	Bibron's Stiletto Snake	Least Concern	4
<i>Lamprophiidae</i>	<i>Atractaspis duerdeni</i>	Duerden's Stiletto Snake	Least Concern	1
<i>Lamprophiidae</i>	<i>Boaedon capensis</i>	Brown House Snake	Least Concern	9
<i>Lamprophiidae</i>	<i>Lycophidion capense capense</i>	Cape Wolf Snake	Least Concern	4
<i>Lamprophiidae</i>	<i>Prosymna sundevallii</i>	Sundevall's Shovel-snout	Least Concern	6
<i>Lamprophiidae</i>	<i>Psammophis brevirostris</i>	Short-snouted Grass Snake	Least Concern	9
<i>Lamprophiidae</i>	<i>Psammophis notostictus</i>	Karoo Sand Snake	Least Concern	1
<i>Lamprophiidae</i>	<i>Psammophis trinasalis</i>	Fork-marked Sand Snake	Least Concern	10
<i>Lamprophiidae</i>	<i>Pseudaspis cana</i>	Mole Snake	Least Concern	7
<i>Lamprophiidae</i>	<i>Xenocalamus bicolor bicolor</i>	Bicoloured Quill-snouted Snake	Least Concern	1
<i>Leptotyphlopidae</i>	<i>Leptotyphlops scutifrons scutifrons</i>	Peters' Thread Snake		6
<i>Pelomedusidae</i>	<i>Pelomedusa subrufa</i>	Central Marsh Terrapin	Least Concern	4
<i>Pythonidae</i>	<i>Python natalensis</i>	Southern African Python	Least Concern	1
<i>Scincidae</i>	<i>Acontias kgalagadi kgalagadi</i>	Striped Blind Legless Skink	Least Concern	6
<i>Scincidae</i>	<i>Panaspis wahlbergi</i>	Wahlberg's Snake-eyed Skink	Least Concern	1
<i>Scincidae</i>	<i>Trachylepis occidentalis</i>	Western Three-striped Skink	Least Concern	12
<i>Scincidae</i>	<i>Trachylepis punctatissima</i>	Speckled Rock Skink	Least Concern	12
<i>Scincidae</i>	<i>Trachylepis punctulata</i>	Speckled Sand Skink	Least Concern	1
<i>Scincidae</i>	<i>Trachylepis spilogaster</i>	Kalahari Tree Skink	Least Concern	38
<i>Scincidae</i>	<i>Trachylepis sulcata sulcata</i>	Western Rock Skink	Least Concern	15
<i>Scincidae</i>	<i>Trachylepis variegata</i>	Variiegated Skink	Least Concern	49
<i>Testudinidae</i>	<i>Psammobates oculifer</i>	Serrated Tent Tortoise	Least Concern	10
<i>Testudinidae</i>	<i>Stigmochelys pardalis</i>	Leopard Tortoise	Least Concern	3
<i>Typhlopidae</i>	<i>Rhinotyphlops lalandei</i>	Delalande's Beaked Blind Snake	Least Concern	1
<i>Varanidae</i>	<i>Varanus albigularis albigularis</i>	Rock Monitor	Least Concern	13
<i>Viperidae</i>	<i>Bitis arietans arietans</i>	Puff Adder	Least Concern	10

11 ANNEX 3. LIST OF AMPHIBIANS

List of amphibians which are likely to occur in the vicinity of the Hotazel 2 site, according to the Southern African Atlas of Frogs. Conservation is from Minter et al. (2004).

Family	Genus	Species	Common name	Red list category
<i>Brevicipitidae</i>	<i>Breviceps</i>	<i>adpersus</i>	Bushveld Rain Frog	Least Concern
<i>Bufo</i>	<i>Amietophrynus</i>	<i>gutturalis</i>	Guttural Toad	Least Concern
<i>Bufo</i>	<i>Amietophrynus</i>	<i>poweri</i>	Power's Toad	Least Concern
<i>Bufo</i>	<i>Amietophrynus</i>	<i>rangeri</i>	Raucous Toad	Least Concern
<i>Bufo</i>	<i>Poyntonophrynus</i>	<i>vertebralis</i>	Southern Pygmy Toad	Least Concern
<i>Bufo</i>	<i>Vandijkophrynus</i>	<i>gariensis</i>	Karoo Toad	Least Concern
<i>Pipidae</i>	<i>Xenopus</i>	<i>laevis</i>	Common Platanna	Least Concern
<i>Pyxicephalidae</i>	<i>Amietia</i>	<i>angolensis</i>	Common or Angola River Frog	Least Concern
<i>Pyxicephalidae</i>	<i>Cacosternum</i>	<i>boettgeri</i>	Common Caco	Least Concern
<i>Pyxicephalidae</i>	<i>Pyxicephalus</i>	<i>adpersus</i>	Giant Bull Frog	Near Threatened
<i>Pyxicephalidae</i>	<i>Tomopterna</i>	<i>cryptotis</i>	Tremelo Sand Frog	Least Concern

12 ANNEX 3. RESULTS OF DEA SCREENING TOOL

A summary of the outputs of the DEA Screening Tool for the site and grid connection corridors is provided below. There are no sensitive features mapped within the study area.


Animal Species Theme

The relative animal species theme sensitivity is provided below and indicates that the site does not fall within the known or modelled range of any terrestrial animal species of high conservation concern. There is a small high sensitivity area around Hotazel, but this is due to the potential presence of Verreaux's Eagle which is outside the scope of this study.



Plant Species Theme

The relative plant species theme sensitivity is provided below and indicates that the site falls within an area considered to be low sensitivity with no known species of conservation concern.



Terrestrial Biodiversity Theme Sensitivity

The terrestrial biodiversity theme sensitivity map is illustrated below. There are no sensitive features within or near the development site.

