

# Antibacterial Ellagic Acid Derivatives and Other Constituents from *Pancovia pedicellaris*

René F. Soh<sup>a</sup>, Jean K. Bankeu<sup>a</sup>, Bruno N. Lenta<sup>b,c</sup>, Brice M. Mba'ning<sup>a</sup>, Silvère Ngouela<sup>a</sup>, Etienne Tsamo<sup>a</sup>, and Norbert Sewald<sup>c</sup>

<sup>a</sup> Department of Organic Chemistry, Faculty of Science, TWAS Research Unit (TRU) of the University of Yaoundé I, P. O. Box 812, Yaoundé, Cameroon

<sup>b</sup> Department of Chemistry, Higher Teacher's Training College, University of Yaoundé I, P. O. Box 47, Yaoundé, Cameroon

<sup>c</sup> Department of Chemistry, Organic and Bioorganic Chemistry, Bielefeld University, P. O. Box 100131, 33501 Bielefeld, Germany

Reprint requests to Prof. Etienne Tsamo. E-mail: [tsamo@cm.refer.org](mailto:tsamo@cm.refer.org) or Silvère Ngouela. E-mail: [sngouela@yahoo.fr](mailto:sngouela@yahoo.fr)

*Z. Naturforsch.* **2009**, *64b*, 1070–1076; received June 16, 2009

Two new ellagic acid derivatives, named panconosides A (**1**) and B (**2**) were isolated from *Pancovia pedicellaris* together with eleven known compounds (**3**–**13**). The structures of **1** and **2**, as well as those of the known compounds were established by spectroscopic methods and by comparison with previously reported data. Compounds **1** and **2** were tested *in vitro* for their antibacterial potential against six strains of microorganisms: *Micrococcus luteus*, *Streptococcus ferus*, *Streptococcus minor*, *Escherichia coli*, *Bacillus subtilis*, and *Pseudomonas agarici*. They were found to exhibit moderate antibacterial activity against all the tested strains compared to standard drugs.

**Key words:** *Pancovia pedicellaris*, Sapindaceae, Ellagic Acid Derivatives, Panconosides A and B, Antibacterial Activity