Funtumia africana

Indigenous

Trade names: Common names:	Bastard wild rubber English: Bush rubber, False rubber Luganda: Nkago Lusoga: Nkago Rukiga: Munyamagosi Runyankore: Munyumatuga, nyamukago Runyoro: Musanda.
Ecology:	The false rubber tree is widespread throughout much of tropical Africa in both evergreen and semi-deciduous forest. In Uganda it is commonly found in the middle storey of rain forests and is common in the Lake Victoria forest zone.
Uses:	Firewood, charcoal (poor quality), timber (for fascia boards, furniture).
Description:	An evergreen tree usually 8-25 m, the trunk cylindrical (only slightly buttressed in large trees) to a tall , narrow but dense crown , the leaves crowded on short dark branchlets. BARK: smooth and thin, brown-black, granular when older, fine vertical fissures. A thin chalk-white latex exudes when cut which remains sticky on the fingers. LEAVES: opposite, leathery dark green above , broadly oval , usually 18-20 cm long , edge wavy, tip slightly pointed, a stalk about 1 cm. Below 8-14 pairs lateral veins with a few hairs at the angles with the midrib (no pits). FLOWERS: yellow-white , fragrant and fleshy , about 1.5 cm across, the tube thick, mouth narrow, the petal lobes longer than the tube and overlapping to the right . FRUIT: a pair of grey-brown , woody flat follicles , sharp-tipped , usually 9-15 cm long , boat-shaped , standing out at right angles from a common stalk . The fruits split along the upper side setting free very many seeds, each with a tuft of 5 cm long silky hairs at one end aiding wind dispersal.
Propagation: Seed: treatment:	Seedlings, wildings. Fruit may be collected at maturity and left to split open gradually in a cool place. If they split on the tree, the seeds are blown away, not necessary.
storage:	can be stored up to two months, but best sown immediately after collection.
Management: Remarks:	Fast growing. Pollarding. This species differs from <i>Funtumia elastica</i> by having sticky latex which does not roll into balls. Neither does it have pits in the axils of the main lateral veins and the fruit are longer. Grow as a pure stand or intercropped with banana, coffee or cocoa. The wood is white, even-textured and cheap furniture is made out of it.

Funtumia africana

Apocynaceae

