

FAUNAL AND FLORAL DIVERSITY AND HABITAT ASSESSMENTS:

FOR THE PROJECT:

***Kagiso Solar Power Plant (Pty.) Ltd
near Hotazel.***

ON:

***THE REMAINING EXTENT OF THE FARM
KAMEELAAR NO. 315, REGISTRATION DIVISION
KURUMAN, NORTHERN CAPE PROVINCE.***

March 2016

Report prepared by:

ENVIRONMENT RESEARCH CONSULTING

ERC forms part of Benah Con cc

cc registration nr: 2005/044901/23

VAT registration nr: 4880237534

Postal address: PO Box 20640, Noordbrug, 2522

E-mail: albie.erc@gmail.com

Mobile: 082 789 4669

Fax: 086 621 4843

Report Reference: ERC 2016-10

Report authors: A.R. Götze (*Pr.Sci.Nat.*)
L. Kotze

TABLE OF CONTENTS

| | | |
|-------|--|----|
| 1 | SPECIALIST INVESTIGATORS | 5 |
| 2 | PROFESSIONAL DECLARATION | 5 |
| 3 | EXECUTIVE SUMMARY | 7 |
| 4 | INTRODUCTION | 11 |
| 4.1 | Background..... | 11 |
| 4.2 | Terms of Reference & General Requirements..... | 12 |
| 4.3 | Aims of the study | 14 |
| 4.4 | Assumptions and Limitations | 14 |
| 4.5 | General Site Description | 15 |
| 5 | FAUNAL ASSESSMENT | 17 |
| 5.1 | Methodology | 17 |
| 5.2 | Results..... | 18 |
| 5.2.1 | Trap transects (Direct sampling) | 19 |
| 5.2.2 | Walk transects (Indirect sampling) | 20 |
| 5.2.3 | Drive transects (Indirect sampling)..... | 20 |
| 5.2.4 | Desktop Study | 20 |
| 6 | FLORISTIC AND GENERAL HABITAT ASSESSMENT | 21 |
| 6.1 | Methodology | 21 |
| 6.2 | General floristic and habitat information..... | 21 |
| 6.3 | Floristic diversity recorded in the study area..... | 23 |
| 6.4 | Description of Broad Vegetation Units in the Study Area..... | 24 |
| 6.4.1 | VU1: <i>Acacia haematoxylon</i> open woodland | 24 |
| 6.4.2 | VU2: <i>Acacia erioloba</i> open woodland..... | 26 |
| 6.5 | Red Data, Protected and Endemic Plant Species..... | 27 |
| 6.6 | Exotic Plant Species | 29 |
| 7 | THREATENED AND PROTECTED ECOSYSTEMS | 29 |
| 8 | IMPACT ASSESSMENT..... | 30 |
| 8.1 | Assessment of expected impacts and relevant mitigation..... | 30 |
| 8.2 | Assessment of the no-go alternative..... | 40 |
| 8.3 | Monitoring requirements | 40 |
| 9 | CONCLUDING REMARKS | 41 |
| 10 | REFERENCES..... | 42 |
| 10.1 | Literature referred to in this report | 42 |
| 10.2 | Other Literature and Field Guides Consulted | 44 |
| 11 | APPENDIX A: lists of faunal species that may occur in the study area | 45 |
| 12 | APPENDIX B: lists of plant families, genera and species recorded in the study area | 50 |
| 13 | APPENDIX C: Recorded positions of red data or protected species.... | 70 |

LIST OF FIGURES

| | |
|--|----|
| Figure 1: Google earth image indicating the regional setting of the study area | 11 |
| Figure 2: Image indicating the preferred site within the farm portion..... | 12 |
| Figure 3: A local scale map. The white border represents the approximate development footprint (preferred site). The numbered red lines represent small mammal trap transects..... | 17 |

| | |
|---|----|
| Figure 4: The direct surrounds of trap transect 1 | 18 |
| Figure 5: The direct surrounds of trap transect 2. | 19 |
| Figure 6: The direct surrounds of trap transect 3. | 19 |
| Figure 7: Image depicting the two vegetation units recorded in the study area | 24 |
| Figure 8: VU1 – portion of <i>Acacia haematoxylon</i> open woodland with good grass cover. | 25 |
| Figure 9: VU1 – portion of <i>Acacia haematoxylon</i> with high abundance of <i>Elephantorrhiza elephantina</i> | 25 |
| Figure 10: VU2 – portion of <i>Acacia erioloba</i> open woodland with moderate grass cover. | 26 |
| Figure 11: Recorded positions of some protected plant species in the study area | 28 |

LIST OF TABLES

| | |
|--|----|
| Table 5-1: Plausible species richness of the study site | 20 |
| Table 6-1: Summary of plant families, genera and species recorded in the study area | 23 |
| Table 6-2: List of protected plant species recorded in the study area | 27 |
| Table 6-3: Protected tree species frequency, density/ha & number of specimens per VU | 29 |
| Table 8-1: Rating system for the evaluation of impacts related to the proposed development | 30 |
| Table 8-2: Assessment of Impact: Loss of habitat for faunal and floral species | 33 |
| Table 8-3: Assessment of Impact: Loss of indigenous faunal and floral species diversity..... | 36 |
| Table 8-4: Assessment of Impact: Loss of faunal and floral species of conservation significance..... | 38 |
| Table 8-5: A summary of the results from the impact assessments..... | 40 |
| Table 11-1: Mammal species likely to occur on or in close proximity to the study site. Species observed during the study period are included; along with the conservation status of each species (protected statuses have been highlighted) | 45 |
| Table 11-2: Reptile species likely to occur on or in close proximity to the study site. Species observed during the study period are included; along with the conservation status of each species (protected statuses have been highlighted) | 47 |
| Table 11-3: Amphibian species likely to occur on or in close proximity to the study site. Species observed during the study period are included; along with the conservation status of each species (protected statuses have been highlighted) | 48 |
| Table 11-4: Protected butterfly species likely to occur on or in close proximity to the site. | 49 |
| Table 12-1: Plant Families and Genera recorded in the study area | 51 |
| Table 12-2: Woody Species – ANGIOSPERMAE – Dicotyledonae | 54 |
| Table 12-3: Graminoids – ANGIOSPERMAE – Monocotyledonae | 55 |

Table 12-4: Herbaceous Shrubs & Forbs – ANGIOSPERMAE –
Monocotyledonae 56

Table 12-5: Herbaceous Shrubs & Forbs – ANGIOSPERMAE – Dicotyledonae
..... 57

Table 12-6: Species list downloaded from POSA (<http://posa.sanbi.org>) on
March 31, 2016, 2:29 pm for QDS 2723 AA and 2723 AC 61

Table 13-1: Coordinates of some recorded ToPS, red data and protected plant
species..... 70

Table 13-2: Calculations of protected tree density in the study area..... 71

1 SPECIALIST INVESTIGATORS

Primary specialist investigator: Albert R. Götze (Pr.Sci.Nat.)

Highest tertiary qualification: M.Sc. *cum laude* (Phytosociology & Restoration Ecology)

Professional affiliation: SACNASP (reg. no. 400011/08)

Background & expertise:

I have been consulting as a professional ecologist, botanist and soil scientist since 2002. I gained valuable experience in the fields of vegetation classification, various restoration disciplines, faunal trapping, soil surveying and wetland surveying during my post graduate studies and later as fieldwork mentor for post graduate ecology students of the Northwest University, Potchefstroom Campus (2008 - 2014), on occasion for game ranch management students of the Tshwane University of Technology. As independent ecological consultant I have experience in various types of scientific floral and faunal studies in the grassland and savannah in Gauteng, North West, Limpopo, Mpumalanga, Free State, Eastern and Northern Cape. I have also on occasion performed vegetation studies in the KwaZulu-Natal savannah and Indian Ocean Coastal Belt, the Eastern Cape thicket, the Western Cape fynbos, Namaqualand, the Karoo and Swaziland. I have 13 years experience in specialist biodiversity, soil and wetland studies and have performed numerous (at least 95) such studies since 2002. I have authored two and co-authored four scientific papers for various local scientific publications since 2004.

Secondary specialist investigator: Leon Kotze

Highest tertiary qualification: BSc (Conservation Ecology)

Professional affiliation: None

Background & expertise:

I am currently studying towards a M.Sc. in Small Mammal Ecology at the University of Witwatersrand and have captured, handled and recorded individual information for close to 500 small mammals to date. My highest qualification is a B.Sc. in Conservation Ecology (obtained at Stellenbosch University) in which Biome ecology and Zoology were my major subjects.

2 PROFESSIONAL DECLARATION

The specialist investigators responsible for conducting this particular specialist faunal and floral and habitat assessment declare that:

- We consider ourselves bound to the rules and ethics of the South African Council for Natural Scientific Professions (SACNASP).

- At the time of conducting the study and compiling this report we did not have any interest, hidden or otherwise, in the proposed development that this study has reference to, except for financial compensation for work done in our professional capacity.
- Work performed for this study was done in an objective manner. Even if this study results in views and findings that are not favorable to the client/applicant, we will not be affected in any manner by the outcome of any environmental process of which this report may form a part, other than being members of the general public.
- We declare that there are no circumstances that may compromise our objectivity in performing this specialist investigation. We do not necessarily object to or endorse the proposed development, but aim to present facts, findings and recommendations based on relevant professional experience and scientific data.
- We do not have any influence over decisions made by the governing authorities.
- Should we, at any point, consider ourselves to be in conflict with any of the above declarations, we shall formally submit a Notice of Withdrawal to all relevant parties and formally register as an Interested and Affected Party.
- We undertake to disclose all material information in our possession that reasonably has or may have the potential of influencing any decision to be taken with respect to the application by a competent authority to such a relevant authority and the applicant.
- We have expertise and experience in conducting specialist reports relevant to this application, including knowledge of the Act, regulations and any guidelines that have relevance to the proposed activity.
- This document and all information contained herein are and will remain the intellectual property Benah Con cc (Environment Research Consulting) and the specialist investigator(s) responsible for conducting the study. This document, in its entirety or any portion thereof, may not be altered in any manner or form, for any purpose without the specific and written consent of the specialist investigator(s).
- We will comply with the Act, regulations and all other applicable legislation.
- All the particulars furnished in this document are true and correct.
- We realize that a false declaration is an offence in terms of Regulation 71 of NEMA and is punishable in terms of section 24F of the Act.



A.R. Götze (M.Sc.; *Pr.Sci.Nat.*)



Leon Kotze (B.Sc.)

3 EXECUTIVE SUMMARY

Based on the findings of this study it is the opinion of the specialist investigators that from a faunal, floral and general ecological point of view, the proposed development is considered favourably, provided that due care is taken to minimise and properly mitigate all identified impacts.

Introduction & Site Description

This study aims to assess the impact that the development of a Photovoltaic Solar Power Plant, on farmland about 10 km southeast of the Northern Cape town of Hotazel, will have on the faunal and floral diversity within the site concerned (development footprint of approximately 250ha), with special reference to Threatened or Protected Species (ToPS). *Environment Research Consulting (ERC)* was contracted to conduct a biodiversity (faunal & floral) and general habitat assessment of portions of the Remaining Extent of the Farm Kameelaar No. 315, near Hotazel in the Northern Cape Province. This report presents the findings of a once off, summer assessment that was conducted over a three day period from 04 to 06 March 2016.

The site falls within the Eastern Kalahari Bushveld Bioregion of the Savanna Biome (Rutherford et al. 2006). Livestock ranching dominates the immediate surrounds and mining activities are a prominent feature in the region. Topography remains more or less homogeneous throughout the study site with no obvious change in slope. Neither permanent nor semi-permanent water bodies were identified from satellite images or after ground-truthing the sites. The site is structurally diverse with good ground cover for most parts. The site is utilised for grazing on occasion and signs of past veldfires (e.g. some trees only produce basal branches) are visible.

Faunal Assessment

Four small mammal trap lines (live trapping) were placed in three different habitat types on 300 m transects. Additionally, non-invasive walk transects were performed daily, documenting all animal sightings (including spoor and / or scat) in writing or by photographs. Drive transects, within the site, averaged in excess of 5 km per day and near similar distances was covered outside the study site daily.

One Murid species was captured during the study period. Only two of the three transects were successful, with mean trap success = 1.85%, and the min. / max. = 1 / 2. Eighteen non-invasive walk transects were performed and at least an hour was spent inspecting the area surrounding each transect, yet only signs of Common Mole-rat activity was recorded. Drive transects, within the site, averaged in excess of 4 km per day and delivered zero sightings. Two additional mammal species were recorded during this effort.

According to literature research, and considering the bioregion, landscape and habitat characteristics, the plausible species richness of the study site is as indicated in Table A according to a ratio of total species vs. total protected and data deficient species.

Table A: Plausible species richness of the study site

| Faunal type | Ratio = total species : total protected and data deficient species |
|-------------|--|
| Mammals | 42 : 7 |
| Reptiles | 33 : 0 |
| Amphibians | 3 : 0 |
| Butterflies | 1 |

Literature research revealed that no animals were restricted or endemic to the area. Some species listed have a low likelihood of occurring within the site, but are nonetheless listed if their habits, habitat requirements and estimated distribution ranges agree with the study findings. The likelihood of any amphibians occurring on the site is low due to the complete lack of local water bodies. No physical record of the listed butterfly occurring in the site exists, but has been included due to the close proximity of the nearest record (i.e. Hotazel) and its “Data deficient” status. Furthermore, this species is endemic to the region and has habitat preferences corresponding with habitat characteristics of the alternative site.

Floristic & Habitat Assessment

A plotless sampling method was used to record floristic and general habitat data. Plant species observed in the study area during the time of the study were recorded and included in plant species lists. The floristic composition of each of the identified broad vegetation units and/or application area are described and discussed.

According to Mucina & Rutherford (2006) the study area falls in the Kuruman Thornveld vegetation type (SVk9). The habitat characteristics of the study area largely resemble the description given for SVk9 above. The site is a slightly undulating sandy plain with shrubs and tall trees. The soil surface is free of rocks. No clearly defined drainage lines were recorded on the preferred or alternative sites.

204 plant species are recorded on the POSA data base of SANBI for the relevant QDS 2723 AA and 2723 AC. A total of only 102 plant species (from 38 plant families and 77 genera) (Table B) were recorded in the study area during the time of the study and indicates moderate species diversity.

Table B: Summary of plant families, genera and species recorded in the study area

| | Families | Genera | Species |
|---|-----------|-----------|------------|
| ANGIOSPERMAE (seed plants): <i>Monocotyledonae:</i> | 6 | 19 | 27 |
| <i>Dicotyledonae:</i> | 32 | 58 | 75 |
| Total: | 38 | 77 | 102 |

Two broad Vegetation Units (VU's) were recorded and described. The three VU's are:

- VU 1: *Acacia haematoxylon* open woodland
- VU 2: *Acacia erioloba* open woodland

Eight plant species of specific conservation significance were recorded in the study area during the study period. One of these species is listed as a Threatened or Protected Species (ToPS) by the National Environmental Management: Biodiversity Act's (Act No. 10 of 2004) list of ToPS as published in Government Gazette no. 36375 of 16 April 2013 (NEMBA ToPS, 2013). Two are listed by Raimondo *et al* (2009) in the South African Red Data list as "Declining" species. Two trees are included in the protected tree species list as published in the National Forests Act (Act no.84 of 1998) (NFA, 1998), and five of the eight are listed as protected and one as specially protected by the Northern Cape Nature Conservation Act (Act no. 9 of 2009) (NCNCA, 2009).

Due to the high numbers of nationally protected trees (NFA, 1998) (i.e. *Acacia erioloba*, *A. haematoxylon* and *Boscia albitrunca*) the individual positions of these species were not individually geo-referenced during this study. Instead a number of belt transects were conducted in each different VU to determine the density at which these species occur in the study area and just beyond. 14 belt transects of 100 x 40 m (4000 m²) were conducted (7 in VU1 and 7 in VU2). All specimens of these species within the belt transect were counted and noted together with the height of each specimen. Differentiation was made between specimens higher than 2 m (> 2 m) and those shorter than 2 m but not less than 1 m (< 2 m = 1 m). Specimens shorter than 1 m were not counted.

During the study no exotic plant species were recorded.

No ecosystems that are threatened and in need of protection according to the National Environmental Management: Biodiversity Act (Act No. 10 of 2004) was recorded in or in the vicinity of the study area.

Impact Assessment

Based on an impact assessment it is evident that there are three expected impacts on the floral ecology within the study area. Table C summarises the findings indicating the significance of the impact before mitigation and management takes place and the likely impact if mitigation and management takes place. From Table C it is evident that prior to management measures being put in place, the impacts are negative-medium or negative-high level impacts. If effective management takes place, all impacts will be reduced to lower level impacts.

Table C: A summary of the results from the impact assessments

| Impact | Not mitigated / managed | Mitigated / managed |
|---|-------------------------|-------------------------------|
| 1. Loss of habitat for faunal and floral species | negative medium impact | negative low impact |
| 2. Loss of indigenous faunal and floral species diversity | negative medium impact | negative low impact |
| 3. Loss of faunal and floral species of conservation significance | negative high impact | negative medium to low impact |

Due the destructive nature of the proposed development to the floristic diversity occurring in the directly affected area and the direct impact it will also have on the faunal diversity of the area on a local scale, the **no-go alternative** will see the area stay in the current condition. The current impacts exerted on the area from an agricultural point of view (not assessed in this study) will remain and, depending on the management strategies employed by the land owner and natural climatic conditions, the current natural condition may improve or deteriorate in future.

A number of monitoring requirements are listed.

Concluding remarks

The low faunal and moderately low floristic species richness and density recorded would equate to an insignificant impact to the regional diversity of plants, mammals, reptiles and amphibians. Although the number of protected faunal species possibly occurring on or in close proximity to the site is low, these deserve consideration.

When considering the area that was investigated during this study (the site and its direct surrounds), from a faunal, floral and general ecological point of view, it is concluded that the preferred site may be accepted for the proposed development.

4 INTRODUCTION

4.1 Background

Under the National Environmental Management Act (107 of 1998) any development that may cause significant damage to the natural environment is by law required to undergo stringent evaluation with the aim of reducing and mitigating the potential environmental impact (www.eia.org.za). This study aims to assess the impact that the development of a Photovoltaic Solar Power Plant, on farmland about 45 km northwest of the Northern Cape town of Kuruman and about 10 km southeast of Hotazel (Figure 1), will have on the faunal and floral diversity within the site concerned (development footprint of approximately 250ha (Figure 2), with special reference to Threatened or Protected Species (ToPS).

Environment Research Consulting (ERC) was contracted to conduct a biodiversity (faunal & floral) and general habitat assessment of a portion of the remaining portion the farm Kameelaar 315 near Hotazel in the Northern Cape Province. This report presents the findings of a once off, summer assessment that was conducted over a three day period from 04 to 06 March 2016.



Figure 1: Google earth image indicating the regional setting of the study area




Figure 2: Image indicating the preferred site within the farm portion.

4.2 Terms of Reference & General Requirements

The scope of the assessment included the PV Solar Energy Facility and its associated structures and infrastructure (such as the power line and access route). The impacts associated with the power line and access route that run beyond the site are considered to be negligible since the actual footprints of disturbance of the power lines is confined to the pylon bases. Furthermore, the power line and access route are aligned with existing roads as far as possible to avoid any negative environmental impacts.

The following ToR and general requirements were supplied by the client:

Specialists in their field of expertise will consider baseline data and identify and assess impacts according to predefined rating scales – refer to attached method of assessment. Specialists will also suggest optional or essential ways in which to mitigate negative impacts and enhance positive impacts. Further, specialists will, where possible, take into consideration the cumulative effects associated with this and other projects which are either developed or in the process of being developed in the local area.

Specialists' reports must comply with Appendix 6 of GNR982 published under sections 24(5), and 44 of the National Environmental Management Act, 1998 (Act No. 107 of 1998), as amended and whereby the following are to be included:

- The details of:
 - the specialist who prepared the report; and

- the expertise of that specialist to compile a specialist report including a curriculum vitae;
- A declaration that the specialist is independent in a form as may be specified by the competent authority;
- An indication of the scope of, and the purpose for which, the report was prepared;
- The date and season of the site investigation and the relevance of the season to the outcome of the assessment;
- A description of the methodology adopted in preparing the report or carrying out the specialised process; the specific identified sensitivity of the site related to the activity and its associated structures and infrastructure;
- An identification of any areas to be avoided, including buffers;
- A map superimposing the activity including the associated structures and infrastructure on the environmental sensitivities of the site including areas to be avoided, including buffers;
- A description of any assumptions made and any uncertainties or gaps in knowledge;
- A description of the findings and potential implications of such findings on the impact of the proposed activity, including identified alternatives on the environment;
- Any mitigation measures for inclusion in the EMPr;
- Any conditions for inclusion in the environmental authorisation;
- Any monitoring requirements for inclusion in the EMPr or environmental authorisation;
- A reasoned opinion-
 - as to whether the proposed activity or portions thereof should be authorised; and
 - if the opinion is that the proposed activity or portions thereof should be authorised, any avoidance, management and mitigation measures that should be included in the EMPr, and where applicable, the closure plan;
- A description of any consultation process that was undertaken during the course of preparing the specialist report;
- A summary and copies of any comments received during any consultation process and where applicable all responses thereto; and
- Any other information requested by the competent authority.

In addition to the above, specialists were expected to:

- Review Scoping Reports, with specific reference to the Comments and Response Report to familiarize with all relevant issues or concerns relevant to their field of expertise;
- In addition to the impacts listed in the Scoping Report, identify any issue or aspect that needs to be assessed and provide expert opinion on any issue in their field of expertise that they deem necessary in order to avoid potential detrimental impacts;
- Assess the degree and extent of all identified impacts (including cumulative impacts) that the preferred project activity and its proposed alternatives, including that of the no-go alternative, may have;
- Identify and list all legislation and permit requirements that are relevant to the development proposal in context of the study;
- Reference all sources of information and literature consulted; and
- Include an executive summary to the report.

4.3 Aims of the study

- Provide a detailed fauna and flora habitat survey.
- Provide a detailed habitat survey of possible threatened or localised plant species and vertebrates.
- Take count and map the location (and provide coordinates) of any protected species or sensitive habitats found on site.
- Evaluate the conservation importance and significance of the site with special emphasis on the current status of threatened species.
- Record possible host plants or food plants of fauna such as butterflies.
- Conduct a literature investigation of possible species that may occur on site.
- Identify potential ecological impacts on fauna and flora that could occur as a result of the development.
- An assessment of the potential direct and indirect impacts resulting from the proposed development during the construction, operation and decommission phases.
- Make recommendations to reduce or minimise impacts, should the development be approved.
- Comment on plant species that can be utilized socially (medicine, food or other cultural or social purposes).

4.4 Assumptions and Limitations

- It was assumed that 3 trap days would be near sufficient for capturing a representative sample of small mammal diversity within the study site (the optimal period being 4 days) (Avenant & Cavallini 2007). The study area,

however, was too large to sample thoroughly for either mammals or reptiles in the time available. Also, faunal observations during the midday heat were unlikely as temperatures ranged between 30°C and 40°C during the time of the study, evoking most animals to reduce activity and seek shelter. Detection was further constrained by the inherently cryptic and/or evasive nature of most wildlife.

- No attempt was made towards sampling Amphibia, due to the small amount of species possibly occurring on the site (Appendix A, Table 11-3) and the complete absence of permanent water bodies.
- Regarding the faunal species lists (Appendix A), it is important to note that distribution maps are often based on limited ecological knowledge and are thus not consistently reliable in predicting a species' occurrence (Hernandez et al. 2006; Newbold 2010). Furthermore, some uncertainty remains regarding the conservation priority for a great deal of southern African species as not all have been assessed and may classify as "Not evaluated" or "Data deficient".
- As no other insect conservation assessments are available we were limited to assessing only butterfly occurrence. In addition, Mecenero et al. (2013) found that butterfly research is lacking in the region concerned.
- It is assumed that plant species flowering only during specific times of the year could be confused with a very similar species of the same genus.
- Some plant species that emerge and bloom during another time of the year or under very specific circumstances may have been missed entirely.
- Due to the conditions encountered during the time of this study some species (faunal & floral) could only be identified up to genus level.
- All species included in the plant species list (Appendix B) were actually observed and recorded in the study area during the time of the study.
- No scientific data was collected or analyzed for the calculation of ecological veld condition. Any comments or observations made in this regard are based on observations, the expert knowledge and relevant professional experience of the specialist investigators.
- *ERC* reserves the right to amend this report, recommendations and/or conclusions at any stage should any additional or otherwise significant information come to light.

4.5 General Site Description

The study site (S27° 15' 26.10" E23° 01' 20.03", alt. 1100m) is located about 45 km northwest of the Northern Cape town of Kuruman and about 10 km southeast of Hotazel next to the R31 between Kuruman and Hotazel. The site falls within the Eastern Kalahari Bushveld Bioregion of the Savanna Biome, with annual precipitation and temperature averaging 362 mm and 17.8 °C, respectively (Rutherford et al. 2006). Livestock ranching dominates the immediate surrounds and mining activities are a prominent feature in the region (*pers. obs.*). Topography remains homogeneous throughout the study

site with no obvious change in slope. Neither permanent nor semi-permanent water bodies were identified from satellite images or after ground-truthing the sites. The site is structurally diverse with good ground cover, for most parts, and a general lack of rockiness. The site is not permanently utilised for grazing (*pers. comm.* Hendrik Venter) and signs of past veldfires (e.g. some trees only produce basal branches) are visible.

5 FAUNAL ASSESSMENT

5.1 Methodology

Before our initial visit, satellite images (Google Earth) of the site were studied and the different habitat types identified (uniform features from an aerial perspective). Upon arrival the sites were ground-truthed. The small mammal trap transects were then placed, at least one transect per habitat type, and each trap baited with a mixture of peanut-butter, oats, sunflower oil and marmite (Avenant & Cavallini 2007). Transects consisted of 30 traps, placed 10 m apart and were checked every morning at 08h00, again at 15h00 and were re-baited daily. Species, sex and reproductive status were recorded for each animal captured, although only species data has been reported here.

Non-invasive walk transects were performed daily, documenting all animal sightings (including spoor and / or scat) in writing or by photograph. Non-invasive walk transects were done along the small mammal trap transects (Figure 2). The area ahead of the observer was observed attentively, specifically for animals flushed from shelter, and stretched a minimum of 250 m. After each trap check a minimum of 20 minutes was designated to examining the environment around each transect, during which I would frequently investigate the area surrounding me with binoculars.



Figure 3: A local scale map. The white border represents the approximate development footprint (preferred site). The numbered red lines represent small mammal trap transects.

Drive transects were also conducted, twice per day, along the same 5 km route. Driving 20-40 km/h the driver would report any animal observed ahead of the vehicle and the passenger would record any animal seen in a 15 m zone to his side of the vehicle. The area surrounding the study site was also extensively travelled throughout the study period and sampled in a similar fashion. This method of sampling served to record the more conspicuous fauna (e.g. tortoises, large mammals and active snakes).

Species lists (Appendix A, Tables 11-1 to 11-4) were constructed using field guides, Red Data Books and Species Atlases (see 'References') complementarily. Only butterflies were considered in constructing an insect species list as they are the subject of the only existing South African insect conservation assessment. As far as information was available, species habitat requirements were also taken into account to substantiate the likelihood of their occurrence. Hence, veldt condition (i.e. pristine or disturbed), vegetation structure and other habitat characteristics contributed to determining the likelihood of a species' occurrence.

No formal consultation process was conducted as part of this faunal study as it was not deemed necessary at the time of the study.

5.2 Results

The study period lasted 3 days and nights with no less than 8 hours spent on the site per day. Ninety traps (30 traps per transect) were placed in three different habitat types and each transect measured approximately 300 m. Traps were removed following the third evening. In an effort to record landscape elements as well as faunal tracks and signs, extensive notes and photographs were taken throughout this period.



Figure 4: The direct surroundings of trap transect 1.



Figure 5: The direct surrounds of trap transect 2.



Figure 6: The direct surrounds of trap transect 3.

5.2.1 Trap transects (Direct sampling)

One Murid species was captured during the study period (Appendix A, Table 11-1). Only transects 1 and 2 (see Figure 2) were successful, with mean trap success (i.e. average amount of occupied traps per day) = 1.85%, and the min. / max. animals captured on a single day = 1 / 2.

5.2.2 Walk transects (Indirect sampling)

Eighteen non-invasive walk transects were performed and at least one hour was spent inspecting the area surrounding each transect. The mounds of the Common Mole-rat (*Cryptomys hottentotus*) were recorded regularly. No other fauna were observed during this effort.

5.2.3 Drive transects (Indirect sampling)

Drive transects, within the site, averaged in excess of 4 km per day and near similar distances was covered outside the study site daily. Two Herpestid species were recorded: Yellow Mongoose and Suricate (Appendix A, Table 11-1). At least one Suricate colony was identified within the site.

5.2.4 Desktop Study

According to literature research, and considering the bioregion, landscape and habitat characteristics, the plausible species richness of the study site is as indicated in Table 5-1 according to the ratio of total species vs. total protected and data deficient species.

Table 5-1: Plausible species richness of the study site

| Faunal type | Ratio = total species : total protected and data deficient species |
|-------------|--|
| Mammals | 42 : 7 |
| Reptiles | 33 : 0 |
| Amphibians | 3 : 0 |
| Butterflies | 1 |

Literature research revealed that no animals were restricted or endemic to the area. Some species listed, for example Brown Hyaena (*Hyaena brunnea*) due to frequent human activity, have a low likelihood of occurring within the site, but are nonetheless listed if their habits, habitat requirements and estimated distribution ranges agree with the study findings. Greater Kudu (*Tragelaphus strepsiceros*) and Eland (*Taurotragus oryx*) have been included as the livestock fencing would not stop them entering and exiting the site. The likelihood of any amphibians occurring on the site is low due to the complete lack of local water bodies. No physical record of the listed butterfly occurring in the site exists (Appendix A, Table 11-4), but has been included due to the close proximity of the nearest record (i.e. Hotazel) and its “Data deficient” status. Furthermore, the species is endemic to the region and has habitat preferences corresponding with habitat characteristics of the study site.

6 FLORISTIC AND GENERAL HABITAT ASSESSMENT

6.1 Methodology

Prior to visiting the site, a list of species that could potentially occur at the site was downloaded from “Plants of Southern Africa” (POSA) on the South African Biodiversity Institute’s (SANBI) website at <http://posa.sanbi.org>. This list is provided at the quarter degree square (QDS) level of accuracy for the QDS 2723 AA and 2723 AC and included in Appendix B. At this broad scale, the list often includes many species that may not be found at the proposed site. However, any species of conservation concern will be indicated in the list and was researched before the site visit in order to know what species of conservation concern should be looked out for.

A visual reconnaissance of the study area was done before surveying commenced. Different homogenous vegetation units were identified and subsequently surveyed on foot and by vehicle in order to determine the floristic composition of each. The following data was recorded:

- All identifiable indigenous plant species (Appendix B) including red data or specially protected and also exotic plant species in each identified vegetation unit.
- General ecological and habitat data that may assist in the description of the floristic component of the study area.

A plotless sampling method was used to record data. Plant species observed in the study area during the time of the study were recorded and included in the plant species lists (Appendix B). The floristic composition of each of the identified broad vegetation units and/or application area are described and discussed. Plant species identification was done following the checklist of Germishuizen & Meyer (2003). Plant material was collected for identification purposes and where necessary the South African National Biodiversity Institute (SANBI) in Pretoria and other specialists were consulted in order to assist in plant species identification. All collected plant material will, if so requested by them, be donated to the South African National Herbarium of SANBI in Pretoria for inclusion into their extensive collection.

No formal consultation process was conducted as part of this floristic study as it was not deemed necessary at the time of the study.

6.2 General floristic and habitat information

According to Mucina & Rutherford (2006) the study area falls in the Kuruman Thornveld vegetation type (SVk9). The following description of SVk9 has been summarized from Mucina & Rutherford (2006):

Kuruman Thornveld

The Kuruman Thornveld vegetation type (SVk9) occurs in the North-West and Northern Cape Provinces on flats from the Postmasburg and Danielskuil area in the south extending via Kuruman to Tsineng and Dewar in the north. The area receives summer and autumn rainfall (MAP: 300 – 450 mm) and winters

are very dry. Summer temperatures reach an average of 35.9 degrees Celsius on average in January and winter nights average –3.3 degrees Celsius in June.

Geologically some Campbell Group dolomite and chert and mostly younger, superficial Kalahari Group sediments occurs. Locally, rocky pavements are also formed in places. Soils in the area are mostly of the Hutton form and consist of windblown red sand, which vary in depth from shallow to deep. Important land types are Ae, Ai, Ag and Ah.

The landscape and vegetation features of SVk9 consists of flat rocky plains and some sloping hills which support a well developed, closed shrub layer dominated by *Acacia mellifera* subsp. *detinens* and *Tarchonanthus camphoratus* and sometimes a well developed open tree layer dominated by *Acacia erioloba*. Important trees and tall shrubs are *Acacia erioloba*, *A. mellifera* subsp. *detinens*, *A. haematoxylon*, *A. hebeclada* subsp. *hebeclada*, *Boscia albitrunca*, *Grewia flava*, *Gymnosporia buxifolia*, *Lycium hirsutum* and *Tarchonanthus camphoratus*. Low shrubs and herbs that generally occur are *Elephantorrhiza elephantina*, *Dicoma schinzii*, *Monechma divaricatum*, *Gisekia africana*, *Gnidia polycephala*, *Harpagophytum procumbens* subsp. *procumbens*, *Helichrysum zeyheri*, *Hermannia comosa*, *Indigofera daleoides*, *Limeum fenestratum*, *Nolletia ciliaris*, *Pentzia calcarea*, *Plinthus sericeus*, *Seddera capensis*, *Tripteris aghillana* and *Vahlia capensis* subsp. *vulgaris*. Grasses of importance are *Aristida meridionalis*, *A. stipitata* subsp. *stipitata*, *Eragrostis lehmanniana*, *E. echinochloidea* and *Melinis repens*. Biogeographically important species include the small trees and tall shrubs *Acacia haematoxylon*, *A. luederitzii* var. *luederitzii* (Kalahari endemics) and *Terminalia sericea* (southernmost distribution in SA interior), the low shrub *Blepharis marginata*, the herb *Corchorus pinnatipartitus* and grass *Digitaria polyphylla* (all three Griqualand West endemics). One species endemic to SVk9, *Gnaphalium englerianum* also occurs.

The conservation status of SVk9 is Least Threatened. A conservation target of 16% is envisioned by conservation authorities, but to date no portion of SVk9 is statutorily conserved. Only 2% is totally transformed by mainly mining activities and settlements. This vegetation type resembles the description of Acocks' (1953) *Kalahari Thornveld and Shrub Bushveld* (VT 16) and also the description in Low and Rebelo (1996) of *Kalahari Plains Thorn Bushveld* (LR 30).

The habitat characteristics of the study area largely resemble the description given for SVk9 above. The site is a slightly undulating sandy plain with shrubs and tall trees. The soil surface is free of rocks. No clearly defined drainage lines were recorded on the preferred or alternative sites.

204 plant species are recorded on the POSA data base of SANBI for the relevant QDS 2723 AA and 2723 AC and is included in Appendix B, Table 12-6. Keep in mind that this list contains species of at least two or three different vegetation types.

6.3 Floristic diversity recorded in the study area

A total of only 102 plant species (from 38 plant families and 77 genera) (Table 6-1 & Appendix B, Table 12-1) were recorded in the study area during the time of the study and indicates moderate species diversity. The woody layer (trees & shrubs) is represented by 12 woody species and the herbaceous layer is made up of 12 graminoids and 76 herbaceous shrubs, dwarf shrubs, geophytes and other herbs. All of the recorded plant species are indigenous to South Africa. From available literature (Pujol 1988; Pooley, 1998; Schmidt *et al* 2002; Shearing & Van Heerden 1994; Van Wyk *et al* 1997; Van Wyk & Gericke 2003) it was established that at least 35 of the recorded plant species in the study area are used for some or other social activities (medicinal, food/nourishment and/or cultural).

Table 6-1: Summary of plant families, genera and species recorded in the study area

| | Families | Genera | Species |
|--|-----------------|---------------|----------------|
| PTERIDOPHYTA (ferns): | 0 | 0 | 0 |
| GYMNOSPERMAE (Coniferous plants): | 0 | 0 | 0 |
| ANGIOSPERMAE (seed plants): | | | |
| <i>Monocotyledonae:</i> | 6 | 19 | 27 |
| <i>Dicotyledonae:</i> | 32 | 58 | 75 |
| Total: | 38 | 77 | 102 |

During the survey, which was done on foot and by vehicle, only taxa that were identifiable during the time of the study were noted and included in the plant species lists in Appendix B (Tables 12-1 to 12-5). The possibility exists that some plant species that emerge and bloom during another time of the year or under very specific circumstances, or species that are locally rare could have been missed during the survey, but on the other hand, the specialist is convinced that the majority of the species occurring in the study area were identified and recorded. The mentioned species lists contain the plant family name and scientific and common names of all plant species that was observed in the study area during the time of the study. Also included is, where applicable, the status of a species, which provides information on endemism, red data status or exotic status. Information on whether a species is utilized for medicinal, cultural or nutritional uses is also provided in the mentioned species lists.

Appendix B, Table 12-1 presents the diversity of plant families, genera and species recorded in the study area. A check list of plant species recorded during this study is included in Tables 12-2 through 12-5 of Appendix B.

6.4 Description of Broad Vegetation Units in the Study Area

Two broad Vegetation Units (VU's) were recorded and are described in the sections below (Figure 7). The three VU's are:

- VU 1: *Acacia haematoxylon* open woodland
- VU 2: *Acacia erioloba* open woodland




Figure 7: Image depicting the two vegetation units recorded in the study area

6.4.1 VU1: *Acacia haematoxylon* open woodland

This VU (Figures 8 & 9) occurs only on more than half the site on slightly undulating terrain on deep sandy soils with no rocks on the soil surface. The vegetation is dominated by woody shrubs and trees. Ecologically speaking VU1 is in a moderate to good veld condition with many high quality grazing plants available in the habitat. The grass cover is good (Figure 9) to moderate and few signs of bush encroachment were noticed. The suffrutex herb *Elephantorrhiza elephantina* is very high in abundance in certain areas of this VU (Figure 9).

The tree species *Acacia haematoxylon* totally dominates the woody cover in this VU. Other tree species and woody shrubs of significance are *Acacia erioloba*, *A. hebeclada*, *A. mellifera* subsp. *detinens*, *Grewia flava* and *Tarchonanthus camphoratus*. Dominant graminoids are *Schmidtia pappophoroides*, *Stipagrostis uniplumis*, *S. ciliata*, *Centropodia glauca* and *Bulbostylis hispidula*. Herbaceous forbs and shrubs that mostly occur in VU3

are *Elephantorrhiza elephantina*, *Aptosimum albomarginatum*, *Sesamum triphyllum*, *Sphenostylis angustifolia*, *Oxygonum delagoense*, *Gisekia africana*, *Plinthus sericeus*, *Limeum viscosum*, *Requienia sphaerosperma*, *Hermannia tomentosa* and *Indigofera alternans*.



Figure 8: VU1 – portion of *Acacia haematoxylon* open woodland with good grass cover.



Figure 9: VU1 – portion of *Acacia haematoxylon* with high abundance of *Elephantorrhiza elephantina*.

During the time of this study 93 plant species (all indigenous) were recorded in VU1. These included 12 woody species, 14 graminoids and 67 herbaceous and dwarf shrubs and other forbs were recorded. From available literature (Pujol 1988; Pooley 1998; Schmidt *et al* 2002; Shearing & Van Heerden 1994; Van Wyk *et al* 1997; Van Wyk & Gericke 2003), it was established that at least 30 of the plant species recorded in VU1 are to some extent utilized for some or other social activity or use (medicinal, nourishment/food, and/or cultural).

6.4.2 VU2: *Acacia erioloba* open woodland

This VU (Figure 10) occurs on deep sandy with no surface rocks on slightly eastward sloping terrain. The vegetation is dominated by trees and tall shrubs. From an ecological point of view VU2 is in a moderate veld condition due to some overgrazing in the past. Although generally the same grass species were recorded as in VU1, the grass cover is poorer in. Low levels of bush encroachment were observed in VU2. As in VU1 moderate dominance of the herbaceous layer by the suffrutex herb *Elephantorrhiza elephantina* was observed in a large portion of this VU.

Dominant tree species in VU2 include *Acacia erioloba*, *A. hebeclada*, *A. mellifera* subsp. *detinens*, *Grewia flava* and *Lycium cinereum*. Dominant graminoids are *Schmidtia pappophoroides*, *Stipagrostis uniplumis*, *S. ciliata*, *Aristida stipitata* and *Centropodia glauca*. Herbaceous shrubs and forbs include *Elephantorrhiza elephantina*, *Senna italica* subsp. *arachoides*, *Gnidia polycephala*, *Heliotropium ciliatum*, *Merremia verecunda*, *Hoffmannseggia burchellii* and *Limeum viscosum*.



Figure 10: VU2 – portion of *Acacia erioloba* open woodland with moderate grass cover.

During the time of this study 60 plant species (all indigenous) were recorded in VU2. These included eight woody species, eight graminoids and 44 herbaceous and dwarf shrubs and other forbs were recorded. It was established from available literature (Pujol 1988; Pooley 1998; Schmidt *et al* 2002; Shearing & Van Heerden 1994; Van Wyk *et al* 1997; Van Wyk & Gericke 2003), that at least 23 of the plant species recorded in this VU are to some extent utilized for some or other social activity or use (medicinal, nourishment/food, and/or cultural).

6.5 Red Data, Protected and Endemic Plant Species

Eight plant species of specific conservation significance were recorded in the study area during the study period. One of these species is listed as a Threatened or Protected Species (ToPS) by the National Environmental Management: Biodiversity Act's (Act No. 10 of 2004) list of ToPS as published in Government Gazette no. 36375 of 16 April 2013 (NEMBA ToPS, 2013). Two are listed by Raimondo *et al* (2009) in the South African Red Data list as "Declining" species. Two trees are included in the protected tree species list as published in the National Forests Act (Act no.84 of 1998) (NFA, 1998), and five of the eight are listed as protected and one as specially protected by the Northern Cape Nature Conservation Act (Act no. 9 of 2009) (NCNCA, 2009).

Table 6-2 lists the recorded ToPS, Red Data listed and protected species relative to the different vegetation units they were recorded in during the time of this study. In Appendix C, Table 13-1 a list appears with the coordinates of recorded protected plant species in the study area. Figure 11 shows the positions of the recorded specimens in relation to the different studied areas. More specimens of these species, which are not listed in Appendix C, do occur in the study area, but due to time constraints these could not be referenced during this study. **It is strongly advised** that once the exact position of development activities and infrastructure has been planned and finalized that a full population study of each affected area be done to determine the population size and extent of these and possibly other protected species within the study area and the relevant appropriate action is then taken.

Table 6-2: List of protected plant species recorded in the study area

Note: abbreviations used in Table 6-2 are as follows:

D – Declining (Raimondo *et al*, 2009); P(SA) – nationally protected tree species (NFA, 1998); P(NC) – provincially protected species (NCNCA, 2009); SP(NC) – provincially specially protected species (NCNCA, 2009); ToPS - threatened or protected species (NEMBA ToPS, 2013).

| SPECIES NAME | FAMILY | GROWTH FORM | SPECIES STATUS | VU | |
|----------------------------|----------------|----------------|----------------|----|---|
| | | | | 1 | 2 |
| <i>Boophone disticha</i> | AMARYLLIDACEAE | Herb, geophyte | D; P(NC) | X | |
| <i>Acacia erioloba</i> | FABACEAE | Tree | D, P(SA) | X | X |
| <i>Acacia haematoxylon</i> | FABACEAE | Tree | P(SA) | X | X |
| <i>Asclepias aurea</i> | APOCYNACEAE | Geophytic herb | P(NC) | | X |

| SPECIES NAME | FAMILY | GROWTH FORM | SPECIES STATUS | VU | |
|--|---------------------|-----------------------|----------------|----|---|
| | | | | 1 | 2 |
| <i>Euphorbia</i> c.f. <i>braunsii</i> | EUPHORBIACEAE | Succulent dwarf shrub | P(NC) | | X |
| <i>Harpagophytum procumbens</i> subsp. <i>procumbens</i> | PEDALIACEAE | Herb | TOPS, SP(NC) | X | X |
| <i>Nerine laticoma</i> | AMARYLLIDACEAE | Geophyte | P(NC) | X | |
| <i>Ruschia</i> c.f. <i>putterillii</i> | MESEMBRYANTHEMACEAE | Succulent herb | P(NC) | X | |



Figure 11: Recorded positions of some protected plant species in the study area

Note: The numbered labels on Figure 11 correspond to the serial number (S/N) in the first column of Table 13-1 of Appendix C.

Due to the high numbers of nationally protected trees (NFA, 1998) (i.e. *Acacia erioloba*, *A. haematoxylon* and *Boscia albitrunca*) the individual positions of these species were not individually geo-referenced during this study. Instead a number of belt transects were conducted in each different VU to determine the density at which these species occur in the study area and just beyond.

14 belt transects of 100 x 40 m (4000 m²) were conducted in the area (7 in VU1 and 7 in VU2) and only the numbers of the three nationally protected trees were considered. All specimens of these species within the belt transect were counted and noted together with the height of each specimen. Differentiation was made between specimens higher than 2 m (> 2 m) and

those shorter than 2 m but not less than 1 m (< 2 m = 1 m). Specimens shorter than 1 m were not counted. Table 6-3 gives a summary of the results of this survey. In Appendix C, Table 13-2 presents the results in detail.

An example for the interpretation of Table 6-3 is as follows: The total number of specimens of, for example, *Acacia haematoxylon* in VU1 is calculated at 13529. This number of specimens is the sum of the *A. haematoxylon* shrubs (1 to < 2 m) i.e. 12549, and the trees (> 2 m) i.e. 980. The total calculated number of *A. haematoxylon* specimens to occur in the study area (300 ha preferred site) is 14357. To calculate the number of specimens of any one of the species for any given surface area, one will take the surface area (in ha) and multiply it with the average species density/ha of the relevant species and VU.

Table 6-3: Protected tree species frequency, density/ha & number of specimens per VU

| VU | VU area (ha) | Average species frequency (as counted on 4000 m ²) | | | | | |
|------------------------------|--------------|--|-------|-------------|----------------------------|------|--------------|
| | | <i>Acacia erioloba</i> | | | <i>Acacia haematoxylon</i> | | |
| | | 1 to <2 m | >2 m | Total | 1 to <2 m | >2 m | Total |
| 1 | | 3.3 | 2.4 | 5.7 | 27.4 | 2.1 | 29.6 |
| 2 | | 3.6 | 18.6 | 22.1 | 1.9 | 0.9 | 2.7 |
| Average species density / ha | | | | | | | |
| 1 | | 8.21 | 6.07 | 14.29 | 68.57 | 5.36 | 73.93 |
| 2 | | 8.93 | 46.43 | 55.36 | 4.64 | 2.14 | 6.79 |
| Number of specimens per VU | | | | | | | |
| 1 | 183 | 1503 | 1111 | 2614 | 12549 | 980 | 13529 |
| 2 | 122 | 1089 | 5664 | 6754 | 566 | 261 | 828 |
| Total: | | | | 9368 | | | 14357 |

6.6 Exotic Plant Species

No exotic plant species were recorded in the study area during the time of this study.

7 THREATENED AND PROTECTED ECOSYSTEMS

No ecosystems that are threatened and in need of protection according to the National Environmental Management: Biodiversity Act (Act No. 10 of 2004) was recorded in or in the vicinity of the study area.

8 IMPACT ASSESSMENT

8.1 Assessment of expected impacts and relevant mitigation

The five tables in the section below (Tables 8-1 to 8-5) serve to summarise the significance of expected and potential impacts on the faunal, floral and habitat features occurring on or directly adjacent to the study area. A summary of expected construction, operational and decommissioning phase impacts are provided. No significant impacts are expected during the pre-construction phase. Tables 8-2, 8-3 and 8-4 present the descriptions of impacts as well as impact assessments according to the method and rating system described in Table 8-1. In addition, Tables 8-2 to 8-4 also indicates mitigatory and management measures needed to minimise the expected ecological impacts.

Table 8-1: Rating system for the evaluation of impacts related to the proposed development

| NATURE | | |
|--|----------------------------|---|
| A brief description of the impact of environmental parameter being assessed in the context of the project. This criterion includes a brief written statement of the environmental aspect being impacted upon by a particular action or activity. | | |
| GEOGRAPHICAL EXTENT | | |
| This is defined as the area over which the impact will be experienced. | | |
| 1 | Site | The impact will only affect the site. |
| 2 | Local/district | Will affect the local area or district. |
| 3 | Province/region | Will affect the entire province or region. |
| 4 | International and National | Will affect the entire country. |
| PROBABILITY | | |
| This describes the chance of occurrence of an impact. | | |
| 1 | Unlikely | The chance of the impact occurring is extremely low (Less than a 25% chance of occurrence). |
| 2 | Possible | The impact may occur (Between a 25% to 50% chance of occurrence). |
| 3 | Probable | The impact will likely occur (Between a 50% to 75% chance of occurrence). |
| 4 | Definite | Impact will certainly occur (Greater than a 75% chance of occurrence). |
| DURATION | | |
| This describes the duration of the impacts. Duration indicates the lifetime of the impact as a result of the proposed activity. | | |
| 1 | Short term | The impact will either disappear with mitigation or will be mitigated through natural processes in a span shorter than the construction phase (0 – 1 years), or the impact will last for the period of a relatively short construction period and a limited recovery time after construction, thereafter it will be entirely negated (0 – 2 years). |
| 2 | Medium term | The impact will continue or last for some time after the construction phase but will be mitigated by direct human action or by natural processes thereafter (2 – 10 years). |
| 3 | Long term | The impact and its effects will continue or last for the entire |

| | | |
|---|-------------------------------|--|
| | | operational life of the development, but will be mitigated by direct human action or by natural processes thereafter (10 – 30 years). |
| 4 | Permanent | The only class of impact that will be non-transitory. Mitigation either by man or natural process will not occur in such a way or such a time span that the impact can be considered indefinite. |
| INTENSITY/ MAGNITUDE | | |
| Describes the severity of an impact. | | |
| 1 | Low | Impact affects the quality, use and integrity of the system/component in a way that is barely perceptible. |
| 2 | Medium | Impact alters the quality, use and integrity of the system/component but system/component still continues to function in a moderately modified way and maintains general integrity (some impact on integrity). |
| 3 | High | Impact affects the continued viability of the system/component and the quality, use, integrity and functionality of the system or component is severely impaired and may temporarily cease. High costs of rehabilitation and remediation. |
| 4 | Very high | Impact affects the continued viability of the system/component and the quality, use, integrity and functionality of the system or component permanently ceases and is irreversibly impaired. Rehabilitation and remediation often impossible. If possible rehabilitation and remediation often unfeasible due to extremely high costs of rehabilitation and remediation. |
| REVERSIBILITY | | |
| This describes the degree to which an impact can be successfully reversed upon completion of the proposed activity. | | |
| 1 | Completely reversible | The impact is reversible with implementation of minor mitigation measures. |
| 2 | Partly reversible | The impact is partly reversible but more intense mitigation measures are required. |
| 3 | Barely reversible | The impact is unlikely to be reversed even with intense mitigation measures. |
| 4 | Irreversible | The impact is irreversible and no mitigation measures exist. |
| IRREPLACEABLE LOSS OF RESOURCES | | |
| This describes the degree to which resources will be irreplaceably lost as a result of a proposed activity. | | |
| 1 | No loss of resource | The impact will not result in the loss of any resources. |
| 2 | Marginal loss of resource | The impact will result in marginal loss of resources. |
| 3 | Significant loss of resources | The impact will result in significant loss of resources. |
| 4 | Complete loss of resources | The impact is result in a complete loss of all resources. |
| CUMULATIVE EFFECT | | |
| This describes the cumulative effect of the impacts. A cumulative impact is an effect which in itself may not be significant but may become significant if added to other existing or potential impacts emanating from other similar or diverse activities as a result of the project activity in question. | | |
| 1 | Negligible cumulative impact | The impact would result in negligible to no cumulative effects. |
| 2 | Low cumulative impact | The impact would result in insignificant cumulative effects. |
| 3 | Medium cumulative | The impact would result in minor cumulative effects. |

| | impact | |
|---|----------------------------|--|
| 4 | High cumulative impact | The impact would result in significant cumulative effects |
| SIGNIFICANCE | | |
| <p>Significance is determined through a synthesis of impact characteristics. Significance is an indication of the importance of the impact in terms of both physical extent and time scale, and therefore indicates the level of mitigation required. The calculation of the significance of an impact uses the following formula: (Extent + probability + reversibility + irreplaceability + duration + cumulative effect) x magnitude/intensity.</p> <p>The summation of the different criteria will produce a non-weighted value. By multiplying this value with the magnitude/intensity, the resultant value acquires a weighted characteristic which can be measured and assigned a significance rating.</p> | | |
| Points | Impact significance rating | Description |
| 6 to 28 | Negative low impact | The anticipated impact will have negligible negative effects and will require little to no mitigation. |
| 6 to 28 | Positive low impact | The anticipated impact will have minor positive effects. |
| 29 to 50 | Negative medium impact | The anticipated impact will have moderate negative effects and will require moderate mitigation measures. |
| 29 to 50 | Positive medium impact | The anticipated impact will have moderate positive effects. |
| 51 to 73 | Negative high impact | The anticipated impact will have significant effects and will require significant mitigation measures to achieve an acceptable level of impact. |
| 51 to 73 | Positive high impact | The anticipated impact will have significant positive effects. |
| 74 to 96 | Negative very high impact | The anticipated impact will have highly significant effects and are unlikely to be able to be mitigated adequately. These impacts could be considered "fatal flaws". |
| 74 to 96 | Positive very high impact | The anticipated impact will have highly significant positive effects. |

Table 8-2: Assessment of Impact: Loss of habitat for faunal and floral species

| Impact | Construction phase | Operational phase | Decommissioning phase |
|--|--|--|--|
| <p><u>IMPACT 1:</u> Loss of habitat for faunal and floral species.</p> | <p>Site clearing and the removal of vegetation leading to loss of faunal & floral habitat.</p> | <p>Ongoing disturbance of soils with general operational activities leading to altered faunal & floral habitat.</p> | <p>Disturbance of soils as part of demolition activities may alter faunal & floral habitat.</p> |
| | <p>Site clearing and the disturbance of soils leading to increased erosion.</p> | <p>Increased run off from paved areas and access roads causing erosion in adjacent areas; Insufficient maintenance of runoff systems leading to erosion.</p> | <p>Disturbance of soils as part of demolition activities leading to increased erosion; Insufficient aftercare and maintenance leading to erosion.</p> |
| | <p>Compaction of soils by construction vehicles.</p> | <p>Ongoing compaction of soils by maintenance vehicles.</p> | <p>Compaction of soils by construction vehicles as part of demolition and rehabilitation activities.</p> |
| | <p>Movement of construction vehicles impacting on habitat through pollution by noise, fuel, oils, hydraulic fluids, etc.</p> | <p>Continued movement of vehicles in the area impacting on habitat through pollution by noise, fuel, oils, hydraulic fluids, etc.</p> | <p>Movement of construction vehicles as part of demolition and rehabilitation activities impacting on habitat through pollution by noise, fuel, oils, hydraulic fluids, etc.</p> |

| | | | |
|--|---|---|--|
| | Disturbance of the soil will transform the vegetation in the study area and create conditions favorable for the establishment of populations of alien and invader plant species as well as common weeds. | Lack of management of transformed habitat will create favorable conditions for the spread of populations of alien and invader plant species to neighboring natural habitats causing further transformation. | Ineffective rehabilitation of impacted areas and failure to implement a comprehensive alien weed control plan may lead to ongoing loss of habitat. |
| | With the development of any infrastructure the fragmentation of natural habitats can occur with the negative effect that the flow of ecosystem services (seed dispersal, pollination, exchanging of genes from one area to the next, etc.) may be interrupted having a negative long term effect on isolated fragments. | Solar panels trap solar energy, effectively altering the microclimate and habitat beneath them. | |

Impact assessment:

| Geographical Extent | Probability | Duration | Intensity / Magnitude | Reversibility | Irreplaceable loss of resources | Cumulative Effect | Significance |
|---------------------|-------------|----------|-----------------------|---------------|---------------------------------|-------------------|---------------------------------------|
| 2 | 4 | 3 | 3 | 2 | 2 | 2 | 45 (negative medium impact) |

Mitigation of Impact 1:

Injudicious and unnecessary destruction of natural vegetation, other than the footprint area of the proposed development, must be avoided at all cost.

To prevent the erosion of topsoil, management measures may include berms, soil traps, hessian curtains and stormwater diversion away from areas susceptible to erosion. Water control structures should be constructed and well maintained to minimize erosion and to create a favorable habitat for the establishment of vegetation during the operation of the development and after decommissioning and rehabilitation.

Wherever possible, any soil that can serve as a growth medium for plants must be stripped and stockpiled for future landscaping and/or rehabilitation after or during the construction phase and should be used as soon as possible after “harvesting” to ensure that seed sources does not become worthless due to decomposition of the seed over time. It must be ensured that such topsoil stockpiles are located outside of any drainage lines and areas susceptible to erosion or siltation. Stockpiles should also be placed away from areas known to contain hazardous substances such as fuel.

All soils compacted as a result of construction activities falling inside the development footprint areas should be ripped and profiled after the construction phase. Special attention should be paid to alien and invasive control within these areas. Alien and invasive vegetation control should take place throughout all development and decommissioning phases to prevent loss of floral habitat.

Proliferation of alien and invasive species is expected within any disturbed areas. These species should be eradicated and controlled to prevent their spread beyond the development/ decommissioning footprint. Alien plant seed dispersal within the top layers of the soil within footprint areas, that will have an impact on future rehabilitation, has to be controlled. A management plan and proper follow-up strategy for the prevention of the establishment and/or further spread of new populations of such species should be developed and enforced.

Vehicles should be well maintained to prevent oil and other chemically based materials to enter the area. Refueling points should be well managed and if any soils are contaminated, it should be stripped and disposed of at a registered hazardous waste dumping site.

After the construction phase and also during the decommissioning/rehabilitation phase, reseeded of indigenous grasses should be done in between the developed infrastructure and all affected areas to re-establish microclimates and niche habitats. These re-seeded areas should be well maintained during the operational phase. Upon decommissioning, all fencing should be removed to re-establish landscape connectivity.

Table 8-3: Assessment of Impact: Loss of indigenous faunal and floral species diversity

| Impact | | Construction phase | Operational phase | Decommissioning phase | | | |
|---|-------------|---|---|--|---------------------------------|-------------------|---------------------------------------|
| IMPACT 2: Loss of indigenous faunal and floral species diversity. | | Site clearance and removal of vegetation for construction of infrastructure and access roads through natural areas leading to a loss of natural species diversity. | Ongoing edge effects from operating the SPP impacting on natural species diversity. | Disturbance of soils as part of demolition activities and ineffective rehabilitation of impacted areas further impacting on natural species diversity. | | | |
| | | Proliferation of alien species may alter plant community structure. Failure to implement a comprehensive alien weed control plan leading to an increase in alien vegetation encroachment. | An increase in alien species leading to altered plant community structure and composition especially in neighboring habitats. | Ineffective rehabilitation of impacted areas and failure to implement a comprehensive alien weed control plan may lead to ongoing loss of natural species diversity. | | | |
| | | | Erosion and sedimentation as a result of operational activities leading to a loss of natural species diversity. | Continued erosion and sedimentation during closure and decommissioning leading to a loss of natural species diversity. | | | |
| Impact assessment: | | | | | | | |
| Geographical Extent | Probability | Duration | Intensity / Magnitude | Reversibility | Irreplaceable loss of resources | Cumulative Effect | Significance |
| 2 | 2 | 4 | 2 | 4 | 3 | 2 | 34 (negative medium impact) |

Mitigation of Impact 2:

An alien vegetation control plan has to be implemented in order to manage alien plant species occurring within the developed and surrounding area.

Removal of the alien and weed species encountered on the property must take place in order to comply with existing legislation (amendments to the regulations under the Conservation of Agricultural Resources Act, 1983 and Section 28 of the National Environmental Management Act, 1998). Removal of species should take place throughout the construction, operational, closure/decommissioning and rehabilitation/maintenance phases. Care should be taken with the choice of herbicides to ensure that no additional impact and loss of indigenous plant species occurs due to the herbicides used. Proper training should be given to contractors/applicators to avoid spraying indigenous vegetation.

Landscaping with local indigenous species is preferable and could include forage and host plants required by pollinators.

After the construction phase and also during the decommissioning/rehabilitation phase, reseeded of local indigenous plant species should be done in between the developed infrastructure and all affected areas to re-establish plant species diversity, which in turn will create habitat for the return of faunal species, especially small mammals and invertebrates. These re-seeded areas should be well maintained during the operational phase.

To prevent the erosion of topsoil, management measures may include berms, soil traps, hessian curtains and stormwater diversion away from areas susceptible to erosion. Water control structures should be constructed and well maintained to minimize erosion and to create a favorable habitat for the establishment of vegetation during the operation of the development and after decommissioning and rehabilitation.

Table 8-4: Assessment of Impact: Loss of faunal and floral species of conservation significance

| Impact | Construction phase | | Operational phase | | Decommissioning phase | | |
|---|--|----------|--|---------------|---|-------------------|---|
| <p>IMPACT 3: Loss of faunal and floral species of conservation significance.</p> | <p>Site clearance and removal of vegetation leading to a loss of any recorded and unrecorded species of conservation significance such as ToPS, Red Data Listed species, Protected species (nationally and/or provincially), plant species with medicinal or other cultural value.</p> | | <p>An increase in alien plant species leading to loss of species of conservation significance such as ToPS, Red Data Listed species, Protected species (nationally and/or provincially), plant species with medicinal or other cultural value by outcompeting these species.</p> | | <p>Ineffective rehabilitation of exposed and impacted areas and failure to implement a comprehensive alien weed control plan leading to ongoing loss of species of conservation significance.</p> | | |
| | | | <p>Erosion and sedimentation as a result of operational activities leading to a loss of species of conservation significance.</p> | | <p>Continued erosion and sedimentation during closure and decommissioning leading to a loss of species of conservation significance.</p> | | |
| Impact assessment: | | | | | | | |
| Geographical Extent | Probability | Duration | Intensity / Magnitude | Reversibility | Irreplaceable loss of resources | Cumulative Effect | Significance |
| 2 | 4 | 4 | 3 | 4 | 2 | 3 | <p>57 (negative high impact)</p> |

Mitigation of Impact 3:

According to SANBI's Guidelines for Environmental Impact Assessments (<http://redlist.sanbi.org/eiaguidelines.php>), *in situ* conservation of species of conservation significance is vital and is recommended as the only option for conserving species of conservation concern. *Ex situ* conservation, i.e. the removal of a subpopulation from its natural habitat to an artificial environment, a practice often termed "search and rescue", will result in the erosion of the inherent genetic diversity and characteristics of that species and increase its risk of extinction in the wild. Similarly, translocation of subpopulations is an unacceptable conservation measure. Translocations are expensive and rarely successful. Even if they are successful, translocated individuals may harm other species within the receiving environment, the translocated individuals may transmit pathogens and/or parasites, and translocation may result in rapid changes in the species itself.

In spite of the above point, if species of conservation significance, and more specifically plant species, are going to be destroyed due to the construction of the proposed development. It may be recommended that these species, especially geophytes, be located and "rescued" by transplanting specimens into a nursery or other safe site until they can be used during rehabilitation and/or landscaping.

Populations of species of conservation significance (ToPS, Red Data Listed species, Protected species (nationally and/or provincially), plant species with medicinal or other cultural value) occurring outside the areas that will be directly impacted by the proposed development needs to be actively conserved in order to conserve a viable, non-fragmented gene pool of these species in the local area.

If possible, developments that jeopardize any large populations of species of conservation significance should be planned in such a way as to avoid the populations and their habitat.

Any specimens of protected plant species known to occur in the vicinity of the development footprint and may potentially be impacted by the development activities, are to be fenced off for the duration of the activity. If these species fall within the development footprint special authorisation is to be obtained from relevant conservation authorities for such species to be cut, disturbed, damaged or destroyed. Applications for such activities should be made to the responsible official within the relevant Northern Cape Nature Conservation Agency.

Based on the above assessment it is evident that there are three expected impacts on the floral ecology within the study area. Table 8-5 summarises the findings indicating the significance of the impact before management takes place (as described in Tables 8-2 to 8-4) and the likely impact if management and mitigation takes place. From Table 8-5 it is evident that prior to management measures being put in place, the impacts are negative-medium or negative-high level impacts. If effective management takes place, all impacts will be reduced to low level impacts.

Table 8-5: A summary of the results from the impact assessments

| Impact | Not mitigated / managed | Mitigated / managed |
|---|--------------------------------|-------------------------------|
| 1. Loss of habitat for faunal and floral species | negative medium impact | negative low impact |
| 2. Loss of indigenous faunal and floral species diversity | negative medium impact | negative low impact |
| 3. Loss of faunal and floral species of conservation significance | negative high impact | negative medium to low impact |

8.2 Assessment of the no-go alternative

Due the destructive nature of the proposed development to the floristic diversity occurring in the directly affected area and the direct impact it will also have on the faunal diversity of the area on a local scale, the no-go alternative will see the area stay in the current condition. The current impacts exerted on the area from an agricultural point of view (not assessed in this study) will remain and, depending on the management strategies employed by the land owner and natural climatic conditions, the current natural condition may improve or deteriorate in future.

8.3 Monitoring requirements

From a floristic point of view the following should be monitored during all phases of the proposed development:

- Floristic diversity of the development area as well as areas directly adjacent.
- Populations of ToPS, Red Data and other protected plant species on neighbouring properties / areas must be assessed and monitored during all project phases.
- The removal of any ToPS, Red Data and other protected plant species must be well monitored and managed. Authorisation, through a provincial and/or national permitting system, is to be obtained from relevant conservation authorities for such species to be cut, disturbed, damaged or destroyed.

From a faunal point of view the following should be monitored:

- Faunal diversity of the areas directly adjacent to the development area.
- During construction any faunal species caught up in the midst of activities, which can be translocated to neighbouring open areas, such as tortoises, should be handled by trained professionals and strictly monitored.
- During the operational phase, as the floristic habitat recovers, the return of especially small mammals should be promoted as these species play an important role in the natural health of an ecosystem. This process can also be monitored by annual or bi-annual monitoring.

9 CONCLUDING REMARKS

The low faunal and moderately low floristic species richness and density recorded would equate to an insignificant impact to the regional diversity of plants, mammals, reptiles and amphibians. Although the number of protected faunal species possibly occurring on or in close proximity to the site is low, these deserve consideration. It must be stressed that the short study period may affect the generation of a representative sample (see also 'Assumptions and Limitations'). We are nonetheless confident in the sampling methods employed as the methodology was designed with the study limitations in mind.

The loss of topsoil and fragmentation of natural habitats that is virtually unavoidable with any type of development, has a negative impact on the regional ecosystem as it disrupts the natural flow of ecosystem services and affects all fauna and flora that are dependent on those habitats. Linear ridges, water courses, wetlands, drainage lines, etc. are especially sensitive to and easily fragmented. A high conservation value is attributed to the plant communities and faunal assemblages of these areas as they contribute significantly to the biodiversity of a region. Care should be taken not to unnecessarily clear or destroy natural vegetation and where possible the rehabilitation of transformed areas and restoration of degraded natural veld should take place in order to improve the ecological health of the floristic component on the property. Development should therefore be planned in such a way that totally transformed areas are chosen for major developments and natural veld, even if it is already degraded and/or fragmented, is avoided as far as possible. A legitimate and well-designed rehabilitation plan must be set in place before mining commences and be strictly enforced on an on-going basis throughout the life of the mine and thereafter.

When considering the different sites (preferred and alternative sites) that were investigated during this study, from a faunal, floral and general ecological point of view, it is concluded that the preferred site may be accepted for the proposed development.

10 REFERENCES

10.1 Literature referred to in this report

Acocks, J.P.H. 1953. Veld Types of South Africa. *Memoirs of the Botanical Survey of South Africa No. 57*. Department of Agriculture and Water Supply, South Africa.

Avenant, N.L. & Cavallini, P. 2007. Correlating rodent community structure with ecological integrity, Tussen-die-Riviere Nature Reserve, Free State province, South Africa. *Integrative Zoology* 2, pp. 212-219.

Cowling, R.M., Richardson, D.M. & Pierce, S.M. 2004. *Vegetation of Southern Africa*, p. 167. Cambridge University Press, London.

Francis, C.D. & Barber, J.R. 2013. A framework for understanding noise impacts on wildlife: an urgent conservation priority. *Frontiers in Ecology and the Environment* 11, pp. 305-313.

Henderson, L. 2001. Alien weeds and Invasive Plants. Plant Protection Research Institute, Agricultural Research Council. Paarl Printers, Cape Town.

Hernandez, P.A., Graham, C.H., Master, L.L. & Albert, D.L. 2006. The effect of sample size and species characteristics on performance of different species distribution modeling methods. *Ecography* 29, pp. 773-785.

Hoffman, M.T., Todd, S., Ntshona, Z. & Turner, S. 1999. *Land degradation in South Africa*. South African National Botanical Institute, Kirstenbosch.

<http://www.eia.org.za/introduction.html> (accessed 24/03/2016 at 11:00AM).

https://www.dwa.gov.za/orange/low_orange/prieska.htm (accessed 24/03/2016 at 14:00PM).

<http://redlist.sanbi.org/eiaguidelines.php>

Low, A.B. & Rebelo, A.G. 1996. *Vegetation of South Africa, Lesotho and Swaziland*. Department of Environmental Affairs and Tourism, Pretoria.

McCrary M.D., McKernan P.A.F. & Wagner W.D. 1982. *Wildlife interactions at solar one: final report*. Rosemead, California.

Mecenero, S., Ball, J.B., Edge, D.A., Hamer, M.L., Henning, G.A., Krüger, M., Pringle, E.L., Terblanche, R.F. & Williams M.C. 2013. *Conservation Assessment of Butterflies of South Africa, Lesotho and Swaziland: Red List and Atlas*. Safronics and the Animal Demography Unit, University of Cape Town.

Mucina, L. & Rutherford, C. 2006. *The vegetation of South Africa, Lesotho and Swaziland*. South African National Biodiversity Institute. Tien Wah Press, Singapore.

NCNCA, 2009. Northern Cape Nature Conservation Act no. 9 of 2009. *Provincial Gazette no 1566, 19 December 2011*.

NEMBA, 2004. National Environmental Management: Biodiversity Act (Act no. 10 of 2004). *Government Gazette No. 26436, Vol. 467, of 7 June 2004*, Pretoria.

NEMBA, 2014. National Environmental Management: Biodiversity Act (Act no. 10 of 2004): Alien and Invasive Species Lists. *Government Gazette No. 37886, Vol. 599, of 01 August 2014*, Pretoria.

NEMBA TOPS, 2013. National Environmental Management: Biodiversity Act (Act 10 of 2004): Publication of Lists of Species that are Threatened or Protected, Activities that are Prohibited and Exemption from Restrictions. *Government Gazette No. 36375, of 16 April 2013. Notice no. 389 of 2013.*

Newbold, T. 2010. Applications and limitations of museum data for conservation and ecology, with particular attention to species distribution models. *Progress in Physical Geography* 34.

NFA, 1998. List of protected tree species under the National Forests Act of 1998 (Act no.84 of 1998), Dept. of Water affairs and Forestry. *Government Gazette No. 30253, Vol. 817*, Pretoria.

Pooley, E. 1998. A Field Guide to Wild Flowers of Kwazulu-Natal and the Eastern Region. Natal Flora Publications Trust, Durban.

Pujol, J. 1988. The Herbalist Handbook – African Flora Medicinal Plants. NaturAfrica, Jean Pujol Natural Healers Foundation, Durban.

Raimondo, D., von Staden, L., Foden, W., Victor, J.E., Helme, N.A., Turner, R.C., Kamundi, D.A. and Manyama, P.A. 2009. Red List of South African Plants. *Strelitzia* 25. South African National Biodiversity Institute, Pretoria.

Rutherford, M.C., Mucina, L. & Powrie, L.W. 2006. Biomes and Bioregions of Southern Africa. *Strelitzia 19: Vegetation of South Africa, Lesotho & Swaziland*, chapter 3. SANBI, Pretoria.

Schmidt, E., Lötter, M. & McClelland, W. 2002. Trees and Shrubs of Mpumalanga and Kruger National Park. Jacana, Johannesburg.

Schradin, C., Krackow, S., Schubert, M., Keller, C., Schradin, B. & Pillay, N. 2007. Regulation of Activity in Desert-Living Striped Mice: The Importance of Basking. *Ethology* 113, pp. 606-614.

Shepard, D.B., Kuhns, A.R., Dreslik, M.J. & Phillips, C.A. 2008. Roads as barriers to animal movement in fragmented landscapes. *Animal Conservation* 11, pp. 288-296.

Shearing, D. & Van Heerden, K. 1994. Karoo. South African Wild Flower Guide 6. Botanical Society of South Africa, Kirstenbosch, Claremont.

Turney, D. & Fthenakis, V. 2011. Environmental impacts from the installation and operation of large-scale solar power plants. *Renewable and Sustainable Energy Reviews* 15, pp. 3261-3270.

Van Wyk, Ben-Erik & Gericke, N. 2003. Peoples Plants, a Guide to Useful Plants of Southern Africa, Briza Publications, Pretoria.

Van Wyk, Ben-Erik, Van Oudtshoorn, B. & Gericke, N. 1997. Medicinal Plants of South Africa, Briza Publications, Pretoria.

10.2 Other Literature and Field Guides Consulted

The following were used for desktop studies and identification of faunal and floral species in the field and not necessarily referred to in the text of this document:

- Alexander, G.J. & Marais, J. 2007. *A Guide to Reptiles of Southern Africa*. Struik Nature, Cape Town.
- Bates, M.F., Branch, W.R, Bauer, A.M., Burger, M., Marais, J., Alexander, G.J. & De Villiers, M.S. 2014. *Atlas and Red List of the Reptiles of South Africa, Lesotho and Swaziland*. SANBI, Pretoria.
- Bromilow. C. 1995. *Problem Plants of South Africa*. Briza Publications cc, Arcadia.
- Du Preez, L. & Carruthers, V. 2009. *A Complete Guide to Frogs of Southern Africa*. Struik Nature, Cape Town.
- Friedmann, Y. & Daly, B, 2004. *Red Data Book of the Mammals of South Africa: A Conservation Assessment*. Endangered Wildlife Trust. South Africa.
- Germishuizen, G. & Fabian, A. 1997. *Wild Flowers of Northern South Africa*. Fernwood Press, Cape Town.
- Manning, J. 2003. *SASOL Eerste Veldgids tot Parasitiese en Vleis-etende Plante van Suider Afrika*. Struik Publishers, Cape Town.
- Republic of South Africa, 2007. *Government Gazette No. 29657, No. R 151*, Pretoria.
- Stuart, C. & Stuart, M. 2015. *Stuart's Field Guide to Mammals of Southern Africa*. Struik Nature, Cape Town.
- Van der Walt, R. 2009. *Wild Flowers of the Limpopo Valley*. Retha Van der Walt, Ludwigslust Game Farms, Musina.
- Van Oudtshoorn, F. 1999. *Guide to the grasses of Southern Africa*. Briza Publications, Pretoria.
- Van Rooyen, N., Bezuidenhout, H. & De Kock, E. 2001. *Blomplante van die Kalahari-duineveld*. Ekotrust cc, Lynwood.
- Van Wyk, A.E. & Malan, S.J. 1997. *Field Guide to the Wild Flowers of the Highveld (2ndedn.)*. Struik Publishers, Cape Town.
- Van Wyk, A.E. & Smith, G.F. 2001. *Regions of floristic endemism in southern Africa. A review with emphasis on succulents*. Umdaus Press, Hatfield, Pretoria.
- Van Wyk, B. & Smith, G.F. 1996. *Guide to Aloes of South Africa*. Briza Publications, Pretoria.
- Van Wyk, B. & Van Wyk, P. 1997. *Field guide to Trees of Southern Africa*, Struik Publishers, Cape Town.

11 APPENDIX A: lists of faunal species that may occur in the study area

Table 11-1: Mammal species likely to occur on or in close proximity to the study site. Species **observed during the study period are included; along with the conservation status of each species (**protected** statuses have been highlighted)**

| Order | Family | Common Name | Species Name | Conservation Status |
|---------------|-----------------|----------------------------------|-----------------------------------|---------------------|
| Macroscelidea | Macroscelididae | Round-eared Sengi | <i>Macroscelides proboscideus</i> | Least concern |
| | | Bushveld Sengi | <i>Elephantulus intufi</i> | Data deficient |
| Eulipotyphla | Erinaceidae | Southern African Hedgehog | <i>Atelerix frontalis</i> | Near threatened |
| Pholidota | Manidae | Ground Pangolin | <i>Smutsia temminckii</i> | Vulnerable |
| Lagomorpha | Leporidae | Cape Hare | <i>Lepus capensis</i> | Least concern |
| | | Scrub Hare | <i>Lepus saxatilis</i> | Least concern |
| Rodentia | Sciuridae | Southern African Ground Squirrel | <i>Xerus inauris</i> | Least concern |
| | Pedetidae | Southern African Springhare | <i>Pedetes capensis</i> | Least concern |
| | Bathyergidae | Common Mole-rat | <i>Cryptomys hottentotus</i> | Least concern |
| | Hystriidae | Cape Porcupine | <i>Hystrix africaeaustralis</i> | Least concern |
| | Muridae | Woosnam's Desert Mouse | <i>Zelotomys woosnami</i> | Least concern |
| | | Pouched Mouse | <i>Saccostumus campestris</i> | Least concern |
| | | Grey Climbing Mouse | <i>Dendromus melanotis</i> | Least concern |
| | | Pygmy Hairy-footed Gerbil | <i>Gerbillurus paeba</i> | Least concern |
| | | Bushveld Gerbil | <i>Gerbilliscus leucogaster</i> | Data deficient |
| | | Highveld Gerbil | <i>Gerbilliscus brantsii</i> | Least concern |
| | | Red Veld Rat | <i>Aethomys chrysophilus</i> | Least concern |
| | | Four-striped Grass Mouse | <i>Rhabdomys</i> spp | Least concern |
| | Black-tailed | <i>Thallomys</i> | Least concern | |

| | | | | |
|-----------------|-----------------|-----------------------------|---------------------------------|-----------------|
| | | Tree Rat | <i>nigricauda</i> | concern |
| | | Southern Multimammate Mouse | <i>Mastomys Coucha</i> | Least concern |
| | | Brant's Whistling Rat | <i>Parotomys brantsii</i> | Least concern |
| Carnivora | Canidae | Cape Fox | <i>Vulpes chama</i> | Least concern |
| | | Bat-eared Fox | <i>Otocyon megalotis</i> | Least concern |
| | | Black-backed Jackal | <i>Canis mesomelas</i> | Least concern |
| | Mustelidae | Honey Badger | <i>Mellivora capensis</i> | Near threatened |
| | | African Striped Weasel | <i>Poecilogale albinucha</i> | Data deficient |
| | | Striped Polecat | <i>Ictonyx striatus</i> | Least concern |
| | Herpestidae | Slender Mongoose | <i>Galerella sanguinea</i> | Least concern |
| | | Yellow Mongoose | <i>Cynictis penicillata</i> | Least concern |
| | | Suricate | <i>Suricata suricatta</i> | Least concern |
| | Viverridae | Small-spotted Genet | <i>Genetta genetta</i> | Least concern |
| | Hyaenidae | Brown Hyaena | <i>Hyaena brunnea</i> | Near threatened |
| | | Aardwolf | <i>Proteles cristatus</i> | Least concern |
| | Felidae | African Wild Cat | <i>Felis silvestris cafra</i> | Least concern |
| | | Small Spotted Cat | <i>Felis nigripes</i> | Least concern |
| | | Caracal | <i>Caracal caracal</i> | Least concern |
| | | Leopard | <i>Panthera pardus</i> | Least concern |
| Tubulidentata | Orycteropodidae | Aardvark | <i>Orycteropus afer</i> | Least concern |
| Cetartiodactyla | Bovidae | Common Eland | <i>Taurotragus oryx</i> | Least concern |
| | | Greater Kudu | <i>Tragelaphus strepsiceros</i> | Least concern |
| | | Steenbok | <i>Raphicerus campestris</i> | Least concern |
| | | Common Duiker | <i>Sylvicapra grimmia</i> | Least concern |

Table 11-2: Reptile species likely to occur on or in close proximity to the study site. Species **observed during the study period are included; along with the conservation status of each species (**protected** statuses have been highlighted)**

| Order | Family | Common Name | Species Name | Conservation Status |
|------------|------------------|-----------------------------|--|---------------------|
| Testudines | Testudinidae | Serrated Tent Tortoise | <i>Psammobates oculifer</i> | Least concern |
| | | Leopard Tortoise | <i>Stigmochelys pardalis</i> | Least concern |
| Squamata | Gekkonidae | Common Giant Gecko | <i>Chondrodactylus angulifer angulifer</i> | Least concern |
| | | Kalahari Ground Gecko | <i>Colopus wahlbergii wahlbergii</i> | Least concern |
| | | Cape Gecko | <i>Pachydactylus capensis</i> | Least concern |
| | | Common Barking Gecko | <i>Ptenopus garrulus garrulus</i> | Least concern |
| | Amphisbaenidae | Pestle-tailed Worm Lizard | <i>Dalophia pistillum</i> | Least concern |
| | | Maurice's Worm Lizard | <i>Monopeltis mauricei</i> | Least concern |
| | Lacertidae | Bushveld Lizard | <i>Heliobolus lugubris</i> | Least concern |
| | | Savanna Lizard | <i>Meroles squamulosus</i> | Least concern |
| | | Spotted Desert Lizard | <i>Meroles suborbitalis</i> | Least concern |
| | | Spotted Sandveld Lizard | <i>Nucras intertexta</i> | Least concern |
| | | Namaqua Sand Lizard | <i>Pedioplanis namaquensis</i> | Least concern |
| | Scincidae | Kgalagadi Legless Skink | <i>Acontias kgalagadi kgalagadi</i> | Least concern |
| | | Western Three-striped Skink | <i>Trachylepis occidentalis</i> | Least concern |
| | | Speckled Sand Skink | <i>Trachylepis punctulata</i> | Least concern |
| | | Kalahari Tree Skink | <i>Trachylepis spilogaster</i> | Least concern |
| | Chamaeleonidae | Common Flap-neck Chameleon | <i>Chamaeleo dilepis dilepis</i> | Least concern |
| | Agamidae | Western Ground Agama | <i>Agama aculeata aculeata</i> | Least concern |
| | Leptotyphlopidae | Peter's Thread | <i>Leptotyphlops</i> | Least concern |

| | | | | |
|--|---------------|--------------------------------|---|-------------------------------------|
| | | Snake | <i>scutifrons</i> | concern |
| | Viperidae | Puff Adder | <i>Bitis arietans arietans</i> | Least concern |
| | Lamprophiidae | Bibron's Stiletto Snake | <i>Atractaspis bibronii</i> | Least concern |
| | | Duerden's Stiletto Snake | <i>Atractaspis duerdeni</i> | Least concern |
| | | Bicoloured Quill-snouted Snake | <i>Xenocalamus bicolor bicolor</i> | Least concern |
| | | Common House Snake | <i>Boaedon capensis</i> | Least concern |
| | | Cape Wolf Snake | <i>Lycophidion capense capense</i> | Least concern |
| | | Fork-marked Sand Snake | <i>Psammophis trinasalis</i> | Least concern |
| | | Sundevall's Shovel-snout | <i>Prosymna sundevalli</i> | Least concern |
| | | Mole Snake | <i>Pseudaspis cana</i> | Least concern |
| | | Elapidae | Common Shield Cobra | <i>Aspidelaps scutatus scutatus</i> |
| | Cape Cobra | | <i>Naja nivea</i> | Least concern |
| | Colubridae | Boomslang | <i>Dispholidus typus</i> | Least concern |
| | | Eastern Tiger Snake | <i>Telescopus semiannulatus semiannulatus</i> | Least concern |

Table 11-3: Amphibian species likely to occur on or in close proximity to the study site. Species **observed during the study period are included; along with the conservation status of each species (**protected** statuses have been highlighted)**

| Order | Family | Common Name | Species Name | Conservation Status |
|-------|----------------|--------------------|-----------------------------|---------------------|
| Anura | Brevicipitidae | Bushveld Rain Frog | <i>Breviceps adspersus</i> | Least concern |
| | | Tremolo Sand Frog | <i>Tomopterna cryptotis</i> | Least concern |
| | | Tandy's Sand Frog | <i>Tomopterna tandyi</i> | Least concern |

Table 11-4: Protected butterfly species likely to occur on or in close proximity to the site.

| Order | Family | Common Name | Species Name | Conservation Status |
|-------------|------------|------------------|---------------------------------|---------------------|
| Lepidoptera | Lycaenidae | Griqua Black Pie | <i>Tuxentius melaena griqua</i> | Data deficient |

12 APPENDIX B: lists of plant families, genera and species recorded in the study area

INDEX

| | |
|--|-------|
| Table 12-1: Plant Families & Genera recorded in the study area | p. 51 |
| Table 12-2: Woody Species – ANGIOSPERMAE – Dicotyledonae..... | p. 54 |
| Table 12-3: Graminoids – ANGIOSPERMAE – Monocotyledonae..... | p. 55 |
| Table 12-4: Herbaceous Shrubs & Forbs (Herbs) – Monocotyledonae..... | p. 56 |
| Table 12-5: Herbaceous Shrubs & Forbs (Herbs) – Dicotyledonae..... | p. 57 |
| Table 12-6: Species list downloaded from POSA (http://posa.sanbi.org) on March 31, 2016, 2:29 pm for QDS 2723 AA and 2723 AC..... | p. 61 |

Abbreviations used in Tables 12-2 to 12-5 are declared as follows:

Under the column SPECIES STATUS:

| | |
|--------|---|
| ToPS | Threatened or Protected Species (NEMBA ToPS, 2013) |
| D | Red data – Declining (Raimondo et al, 2009) |
| P(SA) | Protected nationally (NFA, 1998) |
| P(NC) | Protected in Northern Cape Province (NCNCA, 2009) |
| SP(NC) | Specially Protected in Northern Cape Province (NCNCA, 2009) |

Under the column SOCIAL USE:

| | | |
|---|---|------------------|
| F | – | Food/nourishment |
| M | – | Medicinal |
| C | – | Cultural |

Table 12-1: Plant Families and Genera recorded in the study area

| FAMILY | No. of families | No. of genera per family | GENUS | No. of species per genus | No. of species per genus in VU | |
|------------------------|-----------------|--------------------------|----------------------|--------------------------|--------------------------------|-----------|
| | | | | | 1 | 2 |
| ANGIOSPERMAE | | | | | | |
| MONOCOTYLEDONAE | | | | | | |
| AMARYLLIDACEAE | 1 | 2 | <i>Boophone</i> | 1 | 1 | |
| | | | <i>Nerine</i> | 1 | 1 | |
| ASPARAGACEAE | 1 | 1 | <i>Asparagus</i> | 4 | 4 | 1 |
| COMMELINACEAE | 1 | 1 | <i>Commelina</i> | 1 | 1 | 1 |
| CYPERACEAE | 1 | 2 | <i>Bulbostylis</i> | 1 | 1 | 1 |
| | | | <i>Cyperus</i> | 1 | 1 | |
| HYACINTHACEAE | 1 | 3 | <i>Albuca</i> | 1 | | 1 |
| | | | <i>Dipcadi</i> | 4 | 4 | 1 |
| | | | <i>Ledebouria</i> | 1 | | 1 |
| POACEAE | 1 | 10 | <i>Aristida</i> | 1 | 1 | 1 |
| | | | <i>Cenchrus</i> | 1 | 1 | |
| | | | <i>Centropodia</i> | 1 | 1 | 1 |
| | | | <i>Enneapogon</i> | 1 | 1 | 1 |
| | | | <i>Eragrostis</i> | 1 | 1 | 1 |
| | | | <i>Pogonarthria</i> | 1 | 1 | |
| | | | <i>Schmidtia</i> | 1 | 1 | 1 |
| | | | <i>Stipagrostis</i> | 3 | 3 | 2 |
| | | | <i>Tragus</i> | 1 | 1 | |
| | | | <i>Urochloa</i> | 1 | 1 | |
| Sub-Total: | 6 | 19 | | 27 | 25 | 13 |
| DICOTYLEDONAE | | | | | | |
| ACANTHACEAE | 1 | 1 | <i>Blepharis</i> | 1 | 1 | |
| AIZOACEAE | 1 | 1 | <i>Plinthus</i> | 1 | 1 | |
| AMARANTHACEAE | 1 | 1 | <i>Hermbstaedtia</i> | 1 | 1 | 1 |
| APOCYNACEAE | 1 | 1 | <i>Asclepias</i> | 1 | | 1 |
| ASTERACEAE | 1 | 6 | <i>Berkheya</i> | 1 | | 1 |
| | | | <i>Dicoma</i> | 1 | 1 | 1 |
| | | | <i>Felicia</i> | 2 | 2 | |
| | | | <i>Pentzia</i> | 1 | 1 | |
| | | | <i>Senecio</i> | 1 | 1 | |
| | | | <i>Tarchonanthus</i> | 1 | 1 | |
| BIGNONIACEAE | 1 | 1 | <i>Rhigozum</i> | 1 | 1 | 1 |

| FAMILY | No. of families | No. of genera per family | GENUS | No. of species per genus | No. of species per genus in VU | |
|---------------------|-----------------|--------------------------|------------------------|--------------------------|--------------------------------|---|
| | | | | | 1 | 2 |
| BORAGINACEAE | 1 | 2 | <i>Ehretia</i> | 1 | 1 | |
| | | | <i>Heliotropium</i> | 1 | 1 | 1 |
| CONVOLVULACEAE | 1 | 3 | <i>Convolvulus</i> | 1 | 1 | |
| | | | <i>Ipomoea</i> | 1 | 1 | 1 |
| | | | <i>Merremia</i> | 1 | 1 | 1 |
| CUCURBITACEAE | 1 | 4 | <i>Acanthosicyos</i> | 1 | 1 | 1 |
| | | | <i>Citrullus</i> | 1 | 1 | 1 |
| | | | <i>Cucumis</i> | 1 | 1 | |
| | | | <i>Momordica</i> | 1 | | 1 |
| EBENACEAE | 1 | 1 | <i>Diospyros</i> | 1 | 1 | |
| EUPHORBIACEAE | 1 | 1 | <i>Euphorbia</i> | 2 | 2 | 1 |
| FABACEAE | 1 | 11 | <i>Acacia</i> | 4 | 4 | 4 |
| | | | <i>Chamaecrista</i> | 1 | | 1 |
| | | | <i>Cullen</i> | 1 | 1 | |
| | | | <i>Elephantorrhiza</i> | 1 | 1 | 1 |
| | | | <i>Hoffmannseggia</i> | 1 | 1 | 1 |
| | | | <i>Indigofera</i> | 3 | 3 | 2 |
| | | | <i>Requienia</i> | 1 | 1 | |
| | | | <i>Rhynchosia</i> | 1 | | 1 |
| | | | <i>Senna</i> | 1 | 1 | 1 |
| | | | <i>Sphenostylis</i> | 1 | 1 | 1 |
| | | | <i>Tephrosia</i> | 1 | 1 | 1 |
| GISEKIACEAE | 1 | 1 | <i>Gisekia</i> | 1 | 1 | 1 |
| ILLECEBRACEAE | 1 | 1 | <i>Pollichia</i> | 1 | 1 | 1 |
| LAMIACEAE | 1 | 2 | <i>Acrotome</i> | 1 | 1 | 1 |
| | | | <i>Leucas</i> | 1 | 1 | |
| MENISPERMACEAE | 1 | 1 | <i>Antizoma</i> | 1 | 1 | |
| MESEMBRYANTHEMACEAE | 1 | 1 | <i>Ruschia</i> | 1 | 1 | |
| MOLLUGINACEAE | 1 | 1 | <i>Limeum</i> | 3 | 3 | 2 |
| PEDALIACEAE | 1 | 2 | <i>Harpagophytum</i> | 1 | 1 | 1 |
| | | | <i>Sesamum</i> | 1 | 1 | 1 |
| POLYGALACEAE | 1 | 1 | <i>Polygala</i> | 1 | 1 | |
| POLYGONACEAE | 1 | 1 | <i>Oxygonum</i> | 2 | 2 | 1 |
| PORTULACACEAE | 1 | 1 | <i>Talinum</i> | 2 | 1 | 2 |
| RHAMNACEAE | 1 | 1 | <i>Ziziphus</i> | 1 | 1 | 1 |
| SCROPHULARIACEAE | 1 | 2 | <i>Aptosimum</i> | 2 | 2 | |

| FAMILY | No. of families | No. of genera per family | GENUS | No. of species per genus | No. of species per genus in VU | |
|-------------------|-----------------|--------------------------|--------------------|--------------------------|--------------------------------|-----------|
| | | | | | 1 | 2 |
| | | | <i>Peliostomum</i> | 1 | 1 | 1 |
| SOLANACEAE | 1 | 2 | <i>Lycium</i> | 2 | 2 | 1 |
| | | | <i>Solanum</i> | 1 | 1 | 1 |
| STERCULIACEAE | 1 | 2 | <i>Hermannia</i> | 4 | 4 | 2 |
| | | | <i>Melhania</i> | 1 | 1 | |
| THYMELAEACEAE | 1 | 1 | <i>Gnidia</i> | 1 | 1 | 1 |
| TILIACEAE | 1 | 1 | <i>Grewia</i> | 1 | 1 | 1 |
| VERBENACEAE | 1 | 1 | <i>Lantana</i> | 1 | 1 | |
| VIOLACEAE | 1 | 1 | <i>Hybanthus</i> | 1 | 1 | 1 |
| VISCACEAE | 1 | 1 | <i>Viscum</i> | 1 | | 1 |
| ZYGOPHYLLACEAE | 1 | 1 | <i>Tribulus</i> | 2 | 2 | 2 |
| Sub-Total: | 32 | 58 | | 75 | 68 | 47 |
| Total: | 38 | 77 | | 102 | 93 | 60 |

Table 12-2: Woody Species – ANGIOSPERMAE – Dicotyledonae

| FAMILY | SPECIES NAME | GROWTH FORM | COMMON NAME | | SPECIES STATUS | SOCIAL USE | VU | |
|--------------|--|-------------|---------------------------------|-------------------|----------------|------------|----|---|
| | | | AFRIKAANS | ENGLISH | | | 1 | 2 |
| ASTERACEAE | <i>Tarchonanthus camphoratus</i> L. | Tree | Wildekanferbos | Wild camphor bush | | M | X | |
| BIGNONIACEAE | <i>Rhigozum trichotomum</i> Burch. | Tree | Driedoring | | | | X | X |
| BORAGINACEAE | <i>Ehretia rigida</i> (Thunb.) Druce subsp. <i>rigida</i> | Tree | Deurmekaarbos | Puzzle-bush | | F/C | X | |
| EBENACEAE | <i>Diospyros lycioides</i> Desf. | Tree | Bloubos | Bluebush | | M/F/C | X | |
| FABACEAE | <i>Acacia erioloba</i> E.Mey. | Tree | Kameeldoring | Camel Thorn | D, P(SA) | M/F/C | X | X |
| FABACEAE | <i>Acacia haematoxylon</i> Willd. | Tree | Vaalkameeldoring | Grey Camel Thorn | P(SA) | | X | X |
| FABACEAE | <i>Acacia hebeclada</i> DC. subsp. <i>hebeclada</i> | Tree | Trassiedoring | Candle Thorn | | | X | X |
| FABACEAE | <i>Acacia mellifera</i> (Vahl) Benth. subsp. <i>detinens</i> (Burch.) Brenan | Tree | Swarthaak | Black Thorn | | M/C | X | X |
| RHAMNACEAE | <i>Ziziphus mucronata</i> Willd. subsp. <i>mucronata</i> | Tree | Blinkblaar-wag-'n-bietjie | Buffalo-thorn | | M/F/C | X | X |
| SOLANACEAE | <i>Lycium cinereum</i> Thunb. | Shrub | Kleinkriedoring / Slangbessie | Small Honey-thorn | | C | X | |
| SOLANACEAE | <i>Lycium hirsutum</i> Dunal | Shrub | Rivierkareedoring / Wolwedoring | | | | X | X |
| TILIACEAE | <i>Grewia flava</i> DC. | Tree | Fluweelrosyntjie | Velvet Raisin | | F/C | X | X |

Table 12-3: Graminoids – ANGIOSPERMAE – Monocotyledonae

| FAMILY | SPECIES NAME | GROWTH FORM | COMMON NAME | | SPECIES STATUS | SOCIAL USE | VU | |
|------------|--|----------------|------------------------|-----------------------|----------------|------------|----|---|
| | | | AFRIKAANS | ENGLISH | | | 1 | 2 |
| CYPERACEAE | <i>Bulbostylis hispidula</i> (Vahl) R.W.Haines subsp. <i>pyriformis</i> (Lye) R.W.Haines | Herb, cyperoid | | Veld Bulrush | | | X | X |
| CYPERACEAE | <i>Cyperus obtusiflorus</i> Vahl var. <i>obtusiflorus</i> | Herb, cyperoid | Witbiesie | White-flowered Sedge | | | X | |
| POACEAE | <i>Aristida stipitata</i> Hack. | Grass | Langnaaldsteekgras | Long-awned Grass | | | X | X |
| POACEAE | <i>Cenchrus ciliaris</i> L. | Grass | Bloubuffelgras | Foxtail Buffalo Grass | | | X | |
| POACEAE | <i>Centropodia glauca</i> (Nees) Cope | Grass | Gha-gras | Gha Grass | | | X | X |
| POACEAE | <i>Enneapogon cenchroides</i> (Roem. & Schult.) C.Eragrostis Hubb. | Grass | Negenaaldgras | Nine-awned Grass | | | X | X |
| POACEAE | <i>Eragrostis lehmanniana</i> Nees var. <i>lehmanniana</i> | Grass | Knietjiesgras | Lehmann's Love Grass | | C | X | X |
| POACEAE | <i>Pogonarthria squarrosa</i> (Roem. & Schult.) Pilg. | Grass | Sekelgras | Herringbone Grass | | | X | |
| POACEAE | <i>Schmidtia pappophoroides</i> Steud. | Grass | Sandkweek | Sand Quick | | | X | X |
| POACEAE | <i>Stipagrostis ciliata</i> (Desf.) De Winter var. <i>capensis</i> (Trin. & Rupr.) De Winter | Grass | Langbeen-boesmangras | Tall Bushman Grass | | | X | X |
| POACEAE | <i>Stipagrostis obtusa</i> (Delile) Nees | Grass | Kortbeen-boesmangras | Small Bushman Grass | | | X | |
| POACEAE | <i>Stipagrostis uniplumis</i> (Licht.) De Winter var. <i>uniplumis</i> | Grass | Blinkblaar-boesmangras | Silky Bushman Grass | | | X | X |
| POACEAE | <i>Tragus berteronianus</i> Schult. | Grass | Kousklits | Carrot-seed Grass | | | X | |
| POACEAE | <i>Urochloa</i> species | Grass | Bosveldbeesgras | Bushveld Signal Grass | | | X | |

Table 12-4: Herbaceous Shrubs & Forbs – ANGIOSPERMAE – Monocotyledonae

| FAMILY | SPECIES NAME | GROWTH FORM | COMMON NAME | | SPECIES STATUS | SOCIAL USE | VU | |
|----------------|--|------------------|----------------------------|------------------------|----------------|------------|----|---|
| | | | AFRIKAANS | ENGLISH | | | 1 | 2 |
| AMARYLLIDACEAE | <i>Boophone disticha</i> (L.f.) Herb. | Herb, geophyte | Kopseerblom / Gifbol | Poison Bulb | D; P(NC) | M/C | X | |
| AMARYLLIDACEAE | <i>Nerine laticoma</i> (Ker Gawl.) T.Durand & Schinz | Geophyte | | | P(NC) | M | X | |
| ASPARAGACEAE | <i>Asparagus bechuanicus</i> Baker | Herbaceous shrub | | | | | X | |
| ASPARAGACEAE | <i>Asparagus cooperi</i> Baker | Herbaceous shrub | Katbos | | | | X | |
| ASPARAGACEAE | <i>Asparagus nelsii</i> Schinz | Herbaceous shrub | Sandveldkatbos | | | F | X | X |
| ASPARAGACEAE | <i>Asparagus suaveolens</i> Burch. | Herbaceous shrub | Gewone Katbos / Katdoring | Bushveld Asparagus | | M/F/C | X | |
| COMMELINACEAE | <i>Commelina erecta</i> L. | Herb | Blouselblommetjie | Blue Commelina | | | X | X |
| HYACINTHACEAE | <i>Albuca</i> species | Geophyte | Slymuintjie | | | | | X |
| HYACINTHACEAE | <i>Dipcadi</i> c.f. <i>gracillimum</i> Baker | Geophyte | Ouma-se-groottoon | | | | X | |
| HYACINTHACEAE | <i>Dipcadi platyphyllum</i> Baker | Geophyte | Breëblaar-skaambloemmetjie | Crinkle-leaved Dipcadi | | | X | |
| HYACINTHACEAE | <i>Dipcadi</i> species 1 | Geophyte | | | | | X | X |
| HYACINTHACEAE | <i>Dipcadi</i> species 2 | Geophyte | | | | | X | |
| HYACINTHACEAE | <i>Ledebouria undulata</i> (Jacq.) Jessop | Geophyte | | | | M | | X |

Table 12-5: Herbaceous Shrubs & Forbs – ANGIOSPERMAE – Dicotyledonae

| FAMILY | SPECIES NAME | GROWTH FORM | COMMON NAME | | SPECIES STATUS | SOCIAL USE | VU | |
|----------------|--|--------------------|------------------|----------------------------|----------------|------------|----|---|
| | | | AFRIKAANS | ENGLISH | | | 1 | 2 |
| ACANTHACEAE | <i>Blepharis integrifolia</i> (L.f.) E.Mey. ex Schinz | Herb | Rankklits | | | | X | |
| AIZOACEAE | <i>Plinthus sericeus</i> Pax | Dwarf shrub | Sandganna | | | | X | |
| AMARANTHACEAE | <i>Hermbstaedtia linearis</i> Schinz | Herb | | | | | X | X |
| APOCYNACEAE | <i>Asclepias aurea</i> (Schltr.) Schltr. | Geophytic herb | | | P(NC) | | | X |
| ASTERACEAE | <i>Berkheya</i> species | Herb | | | | | | X |
| ASTERACEAE | <i>Dicoma capensis</i> Less. | Herb | Karmedik | | | M | X | X |
| ASTERACEAE | <i>Felicia filifolia</i> (Vent.) Burt Davy subsp. <i>Filifolia</i> | Dwarf shrub | Draaibossie | Needle-leaved Felicia | | | X | |
| ASTERACEAE | <i>Felicia muricata</i> (Thunb.) Nees subsp. <i>muricata</i> | Herb | Bloublommetjie | | | M/C | X | |
| ASTERACEAE | <i>Pentzia calcarea</i> Kies. | Dwarf shrub | Meerkatkaroo | | | | X | |
| ASTERACEAE | <i>Senecio</i> species | Herb | | | | | X | |
| BORAGINACEAE | <i>Heliotropium ciliatum</i> Kaplan | Herb | | | | | X | X |
| CONVOLVULACEAE | <i>Convolvulus ocellatus</i> Hook.f. var. <i>ocellatus</i> | Herb, climber | | | | | X | |
| CONVOLVULACEAE | <i>Ipomoea bolusiana</i> Schinz subsp. <i>bolusiana</i> | Herb / dwarf shrub | | Narrow-leaved Pink Ipomoea | | F | X | X |
| CONVOLVULACEAE | <i>Merremia verecunda</i> Rendle (1) | Herb, climber | | | | | X | X |
| CUCURBITACEAE | <i>Acanthosicyos naudinianus</i> (Sond.) C.Jeffrey | Prostrate herb | Gemsbokkomkommer | Gemsbok Cucumber | | M/F | X | X |
| CUCURBITACEAE | <i>Citrullus lanatus</i> (Thunb.) Matsum. & Nakai | Herb, climber | Karkoer / Tsamma | Tsamma | | F/C | X | X |

| FAMILY | SPECIES NAME | GROWTH FORM | COMMON NAME | | SPECIES STATUS | SOCIAL USE | VU | |
|---------------|---|-----------------------|---------------------------|--------------------------|----------------|------------|----|---|
| | | | AFRIKAANS | ENGLISH | | | 1 | 2 |
| CUCURBITACEAE | <i>Cucumis zeyheri</i> Sond. | Herb, climber | Wildekomkommer | Wild Cucumber | | M/F | X | |
| CUCURBITACEAE | <i>Momordica balsamina</i> L. | Herb, climber | Laloentjie | | | M/F | | X |
| EUPHORBIACEAE | <i>Euphorbia</i> c.f. <i>braunsii</i> N.E.Br. | Succulent dwarf shrub | Vingerpol | | P(NC) | | X | |
| EUPHORBIACEAE | <i>Euphorbia inaequilatera</i> Sond. var. <i>inaequilatera</i> | Herb | Rooi-opslag | Smooth Creeping Milkweed | | | X | X |
| FABACEAE | <i>Chamaecrista mimosoides</i> (L.) Greene | Herb | Boesmanstee | Fishbone Cassia | | | | X |
| FABACEAE | <i>Cullen tomentosum</i> (Thunb.) J.W.Grimes | Herb | Blouklawer / Rivierklawer | Blue Clover | | | X | |
| FABACEAE | <i>Elephantorrhiza elephantina</i> (Burch.) Skeels | Dwarf shrub | Baswortel | Dwarf Elephant-root | | M/C | X | X |
| FABACEAE | <i>Hoffmannseggia burchellii</i> (DC.) Benth. Ex Oliv. subsp. <i>burchellii</i> | Dwarf shrub, herb | | | | | X | X |
| FABACEAE | <i>Indigofera alternans</i> DC. var. <i>alternans</i> | Herb | Skaapertjie / Klipertjie | | | | X | X |
| FABACEAE | <i>Indigofera charlieriana</i> Schinz var. <i>charlieriana</i> | Herb | | | | | X | X |
| FABACEAE | <i>Indigofera daleoides</i> Benth. ex Harv. var. <i>daleoides</i> | Herb | | | | | X | |
| FABACEAE | <i>Requienia sphaerosperma</i> DC. | Herb | | | | | X | |
| FABACEAE | <i>Rhynchosia totta</i> (Thunb.) DC. var. <i>totta</i> | Herb, climber | | | | F | | X |
| FABACEAE | <i>Senna italica</i> Mill. subsp. <i>arachoides</i> (Burch.) Lock | Herb | Elandsertjie | Eland's Pea | | M | X | X |
| FABACEAE | <i>Sphenostylis angustifolia</i> Sond. | Herb | Wilde-ertjie | Wild sweetpea | | M | X | X |
| FABACEAE | <i>Tephrosia purpurea</i> (L.) Pers. | Herb | | Silver Tephrosia | | | X | X |
| GISEKIACEAE | <i>Gisekia africana</i> (Lour.) Kuntze var. <i>africana</i> | Herb | | | | | X | X |
| ILLECEBRACEAE | <i>Pollichia campestris</i> Ait. | Herbaceous shrub | Teesuikerbossie | Waxberry / Barley | | F | X | X |

| FAMILY | SPECIES NAME | GROWTH FORM | COMMON NAME | | SPECIES STATUS | SOCIAL USE | VU | |
|---------------------|--|----------------|-------------------------|------------------------------|----------------|------------|----|---|
| | | | AFRIKAANS | ENGLISH | | | 1 | 2 |
| | | | | Sugar Bush | | | | |
| LAMIACEAE | <i>Acrotome inflata</i> Benth. | Herb | | | | | X | X |
| LAMIACEAE | <i>Leucas capensis</i> (Benth.) Engl. | Herb / shrub | | | | | X | |
| MENISPERMACEAE | <i>Antizoma angustifolia</i> (Burch.) Miers ex Harv. | Herb, climber | | | | | X | |
| MESEMBRYANTHEMACEAE | <i>Ruschia</i> c.f. <i>putterillii</i> (L.Bolus) L.Bolus | Succulent herb | | | P(NC) | | X | |
| MOLLUGINACEAE | <i>Limeum fenestratum</i> (Fenzl) Heimerl var. <i>fenestratum</i> | Herb | | | | | X | X |
| MOLLUGINACEAE | <i>Limeum sulcatum</i> (Klotzsch) Hutch var. <i>sulcatum</i> | Herb | Kloosaarbossie | | | | X | |
| MOLLUGINACEAE | <i>Limeum viscosum</i> (J.Gay) Fenzl subsp. <i>viscosum</i> var. <i>viscosum</i> | Herb | Kloosaarbossie | | | | X | X |
| PEDALIACEAE | <i>Harpagophytum procumbens</i> (Burch.) DC. ex Meisn. subsp. <i>procumbens</i> | Herb | Duiwelsklou / Ghamaghoe | Devil's Claw / Grapple Plant | TOPS, SP(NC) | M/C | X | X |
| PEDALIACEAE | <i>Sesamum triphyllum</i> Welw. ex Asch. var. <i>triphyllum</i> | Herb | Wildesesam | Wild Sesame | | F | X | X |
| POLYGALACEAE | <i>Polygala leptophylla</i> Burch. | Dwarf shrub | | | | M | X | |
| POLYGONACEAE | <i>Oxygonum delagoense</i> Kuntze | Herb | Menssuuring | Salt-of-the-tortoise | | M/F | X | X |
| POLYGONACEAE | <i>Oxygonum</i> c.f. <i>sinuatum</i> (Hochst. & Steud. ex Meisn.) Dammer | Herb | Dubbeltjie | | | | X | |
| PORTULACACEAE | <i>Talinum</i> c.f. <i>arnotii</i> Hook.f. | Succulent herb | | | | M | | X |
| PORTULACACEAE | <i>Talinum crispatum</i> Dinter ex Poelln. | Succulent herb | Wildevygie | | | M/F | X | X |
| SCROPHULARIACEAE | <i>Aptosimum albomarginatum</i> Marloth & Engl. | Dwarf shrub | Koegab | | | | X | |
| SCROPHULARIACEAE | <i>Aptosimum procumbens</i> (Lehm.) Steud. | Herb | Kankerbos | Carpet Flower | | M | X | |

| FAMILY | SPECIES NAME | GROWTH FORM | COMMON NAME | | SPECIES STATUS | SOCIAL USE | VU | |
|------------------|--|------------------|-------------------------------|-----------------------|----------------|------------|----|---|
| | | | AFRIKAANS | ENGLISH | | | 1 | 2 |
| SCROPHULARIACEAE | <i>Peliostomum leucorrhizum</i> E.Mey. ex Benth. | Dwarf shrub | Springbokkos / Karooviooltjie | Veld Violet | | | X | X |
| SOLANACEAE | <i>Solanum supinum</i> Dunal var. <i>supinum</i> | Herb | | | | | X | X |
| STERCULIACEAE | <i>Hermannia coccocarpa</i> (Eckl. & Zeyh.) Kuntze | Herb | | | | | X | |
| STERCULIACEAE | <i>Hermannia modesta</i> (Ehrenb.) Mast. | Herb | | | | | X | X |
| STERCULIACEAE | <i>Hermannia tomentosa</i> (Turcz.) Schinz ex Engl. | Herbaceous shrub | | | | | X | X |
| STERCULIACEAE | <i>Hermannia vestita</i> Thunb. | Herb | Swaelbossie | | | | X | |
| STERCULIACEAE | <i>Melhania acuminata</i> Mast. var. <i>agnosta</i> (K.Schum.) Willd | Herb | | | | | X | |
| THYMELAEACEAE | <i>Gnidia polycephala</i> (C.A.Mey.) Gilg | Herb | Januariebos | | | | X | X |
| VERBENACEAE | <i>Lantana rugosa</i> Thunb. | Herb | Voëlbrandewyn | Birds' Brandy | | M/F/C | X | |
| VIOLACEAE | <i>Hybanthus</i> c.f. <i>densifolius</i> Engl. | Herb | | Lady's Slipper | | | X | X |
| VISCACEAE | <i>Viscum rotundifolium</i> L.f. | Hemi-parasite | Rooibessie / Voëlent | Red-berried Mistletoe | | M/C | | X |
| ZYGOPHYLLACEAE | <i>Tribulus terrestris</i> L. | Herb | Dubbeltjie | Devil's Thorn | | | X | X |
| ZYGOPHYLLACEAE | <i>Tribulus zeyheri</i> Sond. subsp. <i>zeyheri</i> | Herb | Grootblomdubbeltjie | Devil's Thorn | | | X | X |

Table 12-6: Species list downloaded from POSA (<http://posa.sanbi.org>) on March 31, 2016, 2:29 pm for QDS 2723 AA and 2723 AC

| Family | Species | Threat status | SA Endemic | Growth forms | Lifecycle |
|---------------|---|---------------|------------|---------------------------|-----------|
| ACANTHACEAE | Barleria irritans Nees | LC | No | Dwarf shrub, herb | Perennial |
| ACANTHACEAE | Blepharis integrifolia (L.f.) E.Mey. ex Schinz var. integrifolia | LC | No | Herb | Perennial |
| ACANTHACEAE | Glossochilus burchellii Nees | LC | No | Herb | Perennial |
| ACANTHACEAE | Hypoestes forskalii (Vahl) R.Br. | LC | No | Herb | Perennial |
| ACANTHACEAE | Justicia puberula Immelman | LC | No | Dwarf shrub, herb | Perennial |
| ACANTHACEAE | Monechma divaricatum (Nees) C.B.Clarke | LC | No | Shrub, suffrutex | Perennial |
| AIZOACEAE | Plinthus karooicus I.Verd. | LC | No | Dwarf shrub | Perennial |
| AIZOACEAE | Galenia meziana K.Müll. | LC | No | Dwarf shrub | Perennial |
| AIZOACEAE | Plinthus sericeus Pax | LC | No | Dwarf shrub | Perennial |
| AIZOACEAE | Tetragonia calycina Fenzl | LC | No | Dwarf shrub, succulent | Perennial |
| AIZOACEAE | Trianthema parvifolia E.Mey. ex Sond. var. parvifolia | LC | No | Herb, succulent | Annual |
| AMARANTHACEAE | *Gomphrena celosioides Mart. | Not Evaluated | No | Herb | Perennial |
| AMARANTHACEAE | Hermbstaedtia odorata (Burch.) T.Cooke var. odorata | LC | No | Herb | Perennial |
| AMARANTHACEAE | Kyphocarpa angustifolia (Moq.) Lopr. | LC | No | Herb | Annual |
| AMARANTHACEAE | Pupalia lappacea (L.) A.Juss. var. lappacea | LC | No | Herb | Annual |
| AMARANTHACEAE | Sericorema remotiflora (Hook.f.) Lopr. | LC | No | Herb | Perennial |
| AMARANTHACEAE | Sericorema sericea (Schinz) Lopr. | LC | No | Herb | Annual |
| ANACARDIACEAE | Searsia tridactyla (Burch.) Moffett | LC | No | Shrub, tree | Perennial |
| ANACARDIACEAE | Searsia lancea (L.f.) F.A.Barkley | LC | No | Shrub, tree | Perennial |
| APIACEAE | Deverra burchellii (DC.) Eckl. & Zeyh. | LC | No | Shrub | Perennial |
| APIACEAE | Berula thunbergii (DC.) H.Wolff | LC | No | Herb, hydrophyte | Perennial |
| APOCYNACEAE | Microloma armatum (Thunb.) Schltr. var. burchellii (N.E.Br.) Bruyns | LC | No | Dwarf shrub, shrub | Perennial |
| APOCYNACEAE | Raphionacme velutina Schltr. | LC | No | Geophyte, herb, succulent | Perennial |
| ASPARAGACEAE | Asparagus laricinus Burch. | LC | No | Shrub | Perennial |

| Family | Species | Threat status | SA Endemic | Growth forms | Lifecycle |
|-----------------|---|---------------|------------|------------------|-------------------------|
| ASPARAGACEAE | <i>Asparagus retrofractus</i> L. | LC | No | Scrambler, shrub | Perennial |
| ASPARAGACEAE | <i>Asparagus exuvialis</i> Burch. forma <i>exuvialis</i> | Not Evaluated | No | Shrub | Perennial |
| ASPARAGACEAE | <i>Asparagus nelsii</i> Schinz | LC | No | Shrub | Perennial |
| ASPARAGACEAE | <i>Asparagus suaveolens</i> Burch. | LC | No | Shrub | Perennial |
| ASTERACEAE | <i>Dicoma macrocephala</i> DC. | LC | No | Herb | Perennial |
| ASTERACEAE | <i>Dicoma schinzii</i> O.Hoffm. | LC | No | Herb | Perennial |
| ASTERACEAE | <i>Gazania krebsiana</i> Less. subsp. <i>arctotoides</i> (Less.) Roessler | LC | No | Herb | Perennial |
| ASTERACEAE | <i>Geigeria ornativa</i> O.Hoffm. subsp. <i>ornativa</i> | LC | No | Herb | Annual (occ. perennial) |
| ASTERACEAE | <i>Amellus tridactylus</i> DC. subsp. <i>arenarius</i> (S.Moore) Rommel | LC | No | Herb | Annual |
| ASTERACEAE | * <i>Aster squamatus</i> (Spreng.) Hieron. | Not Evaluated | No | Herb | Annual |
| ASTERACEAE | <i>Felicia fascicularis</i> DC. | LC | No | Shrub | Perennial |
| ASTERACEAE | <i>Felicia namaquana</i> (Harv.) Merxm. | LC | No | Herb | Annual |
| ASTERACEAE | <i>Geigeria filifolia</i> Mattf. | LC | No | Herb | Annual (occ. perennial) |
| ASTERACEAE | <i>Kleinia longiflora</i> DC. | LC | No | Shrub, succulent | Perennial |
| ASTERACEAE | <i>Osteospermum muricatum</i> E.Mey. ex DC. subsp. <i>muricatum</i> | LC | No | Herb | Perennial |
| ASTERACEAE | * <i>Pseudognaphalium luteo-album</i> (L.) Hilliard & B.L.Burt | | No | Herb | Annual |
| ASTERACEAE | <i>Pulicaria scabra</i> (Thunb.) Druce | LC | No | Herb | Annual |
| BORAGINACEAE | <i>Heliotropium nelsonii</i> C.H.Wright | LC | No | Herb | Perennial |
| BORAGINACEAE | <i>Heliotropium strigosum</i> Willd. | LC | No | Herb | Annual (occ. perennial) |
| CAMPANULACEAE | <i>Wahlenbergia androsacea</i> A.DC. | LC | No | Herb | Annual |
| CAPPARACEAE | <i>Cleome angustifolia</i> Forssk. subsp. <i>diandra</i> (Burch.) Kers | LC | No | Herb | Perennial |
| CARYOPHYLLACEAE | <i>Pollichia campestris</i> Aiton | LC | No | Herb | Perennial |
| CARYOPHYLLACEAE | * <i>Spergularia media</i> (L.) C.Presl | Not Evaluated | No | Herb | Perennial |
| CELASTRACEAE | <i>Gymnosporia buxifolia</i> (L.) Szyszyl. | LC | No | Shrub, tree | Perennial |

| Family | Species | Threat status | SA Endemic | Growth forms | Lifecycle |
|----------------|---|---------------|------------|---------------------------|-------------------------|
| CELASTRACEAE | <i>Putterlickia saxatilis</i> (Burch.) M.Jordaan | LC | No | Shrub | Perennial |
| CHENOPODIACEAE | <i>Atriplex semibaccata</i> R.Br. var. <i>appendiculata</i> Aellen | LC | No | Dwarf shrub | Annual (occ. perennial) |
| CHENOPODIACEAE | * <i>Chenopodium ambrosioides</i> L. | Not Evaluated | No | Herb | Annual |
| CHENOPODIACEAE | <i>Salsola rabieana</i> I.Verd. | LC | No | Dwarf shrub, shrub | Perennial |
| COLCHICACEAE | <i>Ornithoglossum vulgare</i> B.Nord. | LC | No | Geophyte (fern) | Perennial |
| COMBRETACEAE | <i>Terminalia sericea</i> Burch. ex DC. | LC | No | Tree | Perennial |
| COMMELINACEAE | <i>Commelina livingstonii</i> C.B.Clarke | LC | No | Herb | Perennial |
| CONVOLVULACEAE | <i>Convolvulus ocellatus</i> Hook.f. var. <i>ocellatus</i> | LC | No | Herb | Perennial |
| CONVOLVULACEAE | <i>Ipomoea obscura</i> (L.) Ker Gawl. var. <i>obscura</i> | LC | No | Herb | Perennial |
| CONVOLVULACEAE | <i>Seddera capensis</i> (E.Mey. ex Choisy) Hallier f. | LC | No | Suffrutex | Perennial |
| CONVOLVULACEAE | <i>Xenostegia tridentata</i> (L.) D.F.Austin & Staples subsp. <i>angustifolia</i> (Jacq.) Lejoly & Lisowski | LC | No | Herb | Perennial |
| CONVOLVULACEAE | <i>Evolvulus alsinoides</i> (L.) L. | LC | No | Herb | Annual (occ. perennial) |
| CUCURBITACEAE | <i>Acanthosicyos naudinianus</i> (Sond.) C.Jeffrey | LC | No | Herb, succulent | Perennial |
| CUCURBITACEAE | <i>Corallocarpus triangularis</i> Cogn. | LC | No | Climber, herb, succulent | Perennial |
| CUCURBITACEAE | <i>Cucumis myriocarpus</i> Naudin subsp. <i>myriocarpus</i> | LC | No | Herb | Annual |
| CUCURBITACEAE | <i>Coccinia rehmannii</i> Cogn. | LC | No | Climber, herb, succulent | Perennial |
| CUCURBITACEAE | <i>Trochomeria debilis</i> (Sond.) Hook.f. | LC | No | Climber, herb, succulent | Perennial |
| CYPERACEAE | <i>Bulbostylis burchellii</i> (Ficalho & Hiern) C.B.Clarke | LC | No | Cyperoid, herb, mesophyte | Perennial |
| CYPERACEAE | <i>Bulbostylis hispidula</i> (Vahl) R.W.Haines subsp. <i>pyriformis</i> (Lye) R.W.Haines | LC | No | Cyperoid, herb, mesophyte | Annual |
| CYPERACEAE | <i>Cyperus margaritaceus</i> Vahl var. <i>margaritaceus</i> | LC | No | Cyperoid, herb, mesophyte | Perennial |
| CYPERACEAE | <i>Cyperus marlothii</i> Boeckeler | LC | No | Cyperoid, herb, | Perennial |

| Family | Species | Threat status | SA Endemic | Growth forms | Lifecycle |
|---------------|---|---------------|------------|--|-------------------------|
| | | | | mesophyte | |
| CYPERACEAE | <i>Bolboschoenus maritimus</i> (L.) Palla | LC | No | Cyperoid, emergent hydrophyte, helophyte, herb | Perennial |
| CYPERACEAE | <i>Cyperus longus</i> L. var. <i>tenuiflorus</i> (Rottb.) Boeck. | LC | No | Cyperoid, helophyte, herb | Perennial |
| CYPERACEAE | <i>Scirpoides dioeca</i> (Kunth) Browning | LC | No | Cyperoid, herb, mesophyte | Perennial |
| ELATINACEAE | <i>Bergia anagalloides</i> E.Mey. ex Fenzl | LC | No | Herb | Annual |
| EUPHORBIACEAE | <i>Euphorbia inaequilatera</i> Sond. var. <i>inaequilatera</i> | LC | No | Dwarf shrub, herb | Annual |
| EUPHORBIACEAE | <i>Euphorbia duseimata</i> R.A.Dyer | LC | No | Dwarf shrub, succulent | Perennial |
| EUPHORBIACEAE | <i>Euphorbia pseudotuberosa</i> Pax | LC | No | Dwarf shrub, succulent | Perennial |
| EUPHORBIACEAE | <i>Euphorbia wilmaniae</i> Marloth | LC | No | Dwarf shrub, succulent | Perennial |
| EUPHORBIACEAE | <i>Tragia dioica</i> Sond. | LC | No | Dwarf shrub, herb | Perennial |
| FABACEAE | <i>Elephantorrhiza elephantina</i> (Burch.) Skeels | LC | No | Dwarf shrub, shrub, suffrutex | Annual (occ. perennial) |
| FABACEAE | <i>Indigofera alternans</i> DC. var. <i>alternans</i> | LC | No | Herb | Perennial |
| FABACEAE | <i>Rhynchosia confusa</i> Burt Davy | Not Evaluated | No | Climber, herb | Perennial |
| FABACEAE | <i>Rhynchosia totta</i> (Thunb.) DC. var. <i>totta</i> | LC | No | Climber, herb | Perennial |
| FABACEAE | <i>Rhynchosia venulosa</i> (Hiern) K.Schum. | Not Evaluated | No | Climber, herb | Perennial |
| FABACEAE | <i>Tephrosia burchellii</i> Burt Davy | LC | No | Herb | Annual (occ. perennial) |
| FABACEAE | <i>Tephrosia longipes</i> Meisn. subsp. <i>longipes</i> var. <i>longipes</i> | LC | No | Dwarf shrub, herb, shrub | Annual (occ. perennial) |
| FABACEAE | <i>Tephrosia purpurea</i> (L.) Pers. subsp. <i>leptostachya</i> (DC.) Brummitt var. <i>leptostachya</i> | LC | No | Herb | Annual (occ. perennial) |

| Family | Species | Threat status | SA Endemic | Growth forms | Lifecycle |
|---------------|---|---------------|------------|-----------------------|-------------------------|
| FABACEAE | <i>Acacia karroo</i> Hayne | LC | No | Shrub, tree | Perennial |
| FABACEAE | <i>Crotalaria griquensis</i> L.Bolus | LC | No | Herb | Perennial |
| FABACEAE | <i>Indigofera daleoides</i> Benth. ex Harv. var. <i>daleoides</i> | LC | No | Herb | Perennial |
| FABACEAE | <i>Melolobium microphyllum</i> (L.f.) Eckl. & Zeyh. | LC | No | Dwarf shrub, shrub | Perennial |
| FABACEAE | <i>Otoptera burchellii</i> DC. | LC | No | Climber, herb, shrub | Perennial |
| FABACEAE | <i>Pomaria lactea</i> (Schinz) B.B.Simpson & G.P.Lewis | LC | No | [No lifeform defined] | [No lifecycle defined] |
| GISEKIACEAE | <i>Gisekia africana</i> (Lour.) Kuntze var. <i>pedunculata</i> (Oliv.) Brenan | Not Evaluated | No | Herb | Annual (occ. perennial) |
| GISEKIACEAE | <i>Gisekia pharnacioides</i> L. var. <i>pharnacioides</i> | LC | No | Herb | Annual |
| HYACINTHACEAE | <i>Dipcadi viride</i> (L.) Moench | LC | No | Geophyte | Perennial |
| HYACINTHACEAE | <i>Dipcadi marlothii</i> Engl. | LC | No | Geophyte | Perennial |
| HYACINTHACEAE | <i>Ledebouria apertiflora</i> (Baker) Jessop | LC | No | Geophyte | Perennial |
| IRIDACEAE | <i>Babiana hypogaea</i> Burch. | LC | No | Geophyte, herb | Perennial |
| JUNCACEAE | <i>Juncus rigidus</i> Desf. | LC | No | Helophyte, herb | Perennial |
| LAMIACEAE | <i>Leucas capensis</i> (Benth.) Engl. | LC | No | Dwarf shrub | Perennial |
| LAMIACEAE | <i>Stachys spathulata</i> Burch. ex Benth. | LC | No | Herb | Perennial |
| LAMIACEAE | <i>Ocimum americanum</i> L. var. <i>americanum</i> | LC | No | Herb | Perennial |
| LAMIACEAE | <i>Ocimum filamentosum</i> Forssk. | LC | No | Herb | Perennial |
| LOBELIACEAE | <i>Lobelia thermalis</i> Thunb. | LC | No | Herb | Perennial |
| MALVACEAE | <i>Abutilon austro-africanum</i> Hochr. | LC | No | Dwarf shrub | Perennial |
| MALVACEAE | <i>Grewia flava</i> DC. | LC | No | Shrub | Perennial |
| MALVACEAE | <i>Hibiscus ludwigii</i> Eckl. & Zeyh. | LC | No | Herb, shrub | Perennial |
| MALVACEAE | <i>Hibiscus pusillus</i> Thunb. | LC | No | Herb | Perennial |
| MALVACEAE | <i>Melhania burchellii</i> DC. | LC | No | Herb | Perennial |
| MALVACEAE | <i>Melhania virescens</i> (K.Schum.) K.Schum. | LC | No | Dwarf shrub | Perennial |
| MALVACEAE | <i>Sida chrysantha</i> Ulbr. | LC | No | Dwarf shrub | Perennial |
| MALVACEAE | <i>Sida cordifolia</i> L. subsp. <i>cordifolia</i> | LC | No | Dwarf shrub | Annual (occ. |

| Family | Species | Threat status | SA Endemic | Growth forms | Lifecycle |
|----------------|---|---------------|------------|-----------------------|-------------------------|
| | | | | | perennial) |
| MALVACEAE | <i>Hermannia comosa</i> Burch. ex DC. | LC | No | Herb | Perennial |
| MALVACEAE | <i>Hermannia modesta</i> (Ehrenb.) Mast. | LC | No | Dwarf shrub, herb | Perennial |
| MALVACEAE | <i>Hermannia tomentosa</i> (Turcz.) Schinz ex Engl. | LC | No | Herb | Perennial |
| MALVACEAE | <i>Hibiscus engleri</i> K.Schum. | LC | No | Herb | Perennial |
| MALVACEAE | <i>Hibiscus fleckii</i> Gürke | LC | No | Herb | Perennial |
| MALVACEAE | <i>Hibiscus micranthus</i> L.f. var. <i>micranthus</i> | LC | No | Herb, shrub | Perennial |
| MALVACEAE | <i>Sida ovata</i> Forssk. | LC | No | Dwarf shrub, herb | Annual (occ. perennial) |
| MOLLUGINACEAE | <i>Limeum aethiopicum</i> Burm.f. var. <i>intermedium</i> Friedrich | Not Evaluated | No | [No lifeform defined] | [No lifecycle defined] |
| MOLLUGINACEAE | <i>Limeum viscosum</i> (J.Gay) Fenzl subsp. <i>transvaalense</i> Friedrich | LC | No | Herb | Annual |
| MONTINIACEAE | <i>Montinia caryophyllacea</i> Thunb. | LC | No | Shrub | Perennial |
| PHYLLANTHACEAE | <i>Phyllanthus maderaspatensis</i> L. | LC | No | Herb | Annual (occ. perennial) |
| PHYLLANTHACEAE | <i>Phyllanthus parvulus</i> Sond. var. <i>garipensis</i> (E.Mey. ex Drège) Radcl.-Sm. | LC | No | Dwarf shrub, herb | Perennial |
| PHYLLANTHACEAE | <i>Phyllanthus parvulus</i> Sond. var. <i>parvulus</i> | LC | No | Dwarf shrub, herb | Perennial |
| POACEAE | <i>Andropogon chinensis</i> (Nees) Merr. | LC | No | Graminoid | Perennial |
| POACEAE | <i>Andropogon schirensis</i> Hochst. ex A.Rich. | LC | No | Graminoid | Perennial |
| POACEAE | <i>Anthephora argentea</i> Gooss. | LC | No | Graminoid | Perennial |
| POACEAE | <i>Aristida congesta</i> Roem. & Schult. subsp. <i>congesta</i> | LC | No | Graminoid | Perennial |
| POACEAE | <i>Aristida meridionalis</i> Henrard | LC | No | Graminoid | Perennial |
| POACEAE | <i>Aristida stipitata</i> Hack. subsp. <i>spicata</i> (De Winter) Melderis | LC | No | Graminoid | Perennial |
| POACEAE | <i>Aristida vestita</i> Thunb. | LC | No | Graminoid | Perennial |
| POACEAE | <i>Brachiaria nigropedata</i> (Ficalho & Hiern) Stapf | LC | No | Graminoid | Perennial |
| POACEAE | <i>Cenchrus ciliaris</i> L. | LC | No | Graminoid | Perennial |
| POACEAE | <i>Cymbopogon pospischillii</i> (K.Schum.) C.E.Hubb. | Not Evaluated | No | Graminoid | Perennial |

| Family | Species | Threat status | SA Endemic | Growth forms | Lifecycle |
|---------|---|---------------|------------|--------------|-------------------------|
| POACEAE | <i>Cynodon dactylon</i> (L.) Pers. | LC | No | Graminoid | Perennial |
| POACEAE | <i>Digitaria polyphylla</i> Henrard | LC | No | Graminoid | Perennial |
| POACEAE | <i>Elionurus muticus</i> (Spreng.) Kunth | LC | No | Graminoid | Perennial |
| POACEAE | <i>Enneapogon cenchroides</i> (Licht. ex Roem. & Schult.) C.E.Hubb. | LC | No | Graminoid | Annual (occ. perennial) |
| POACEAE | <i>Enneapogon scoparius</i> Stapf | LC | No | Graminoid | Perennial |

| | | | | | |
|---------|--|---------------|----|-----------|-------------------------|
| POACEAE | * <i>Eragrostis barrelieri</i> Daveau | Not Evaluated | No | Graminoid | Annual |
| POACEAE | <i>Eragrostis biflora</i> Hack. ex Schinz | LC | No | Graminoid | Annual |
| POACEAE | <i>Eragrostis curvula</i> (Schrud.) Nees | LC | No | Graminoid | Perennial |
| POACEAE | <i>Eragrostis lehmanniana</i> Nees var. <i>lehmanniana</i> | LC | No | Graminoid | Perennial |
| POACEAE | <i>Eragrostis nindensis</i> Ficalho & Hiern | LC | No | Graminoid | Perennial |
| POACEAE | <i>Eragrostis pallens</i> Hack. | LC | No | Graminoid | Perennial |
| POACEAE | <i>Eragrostis rigidior</i> Pilg. | LC | No | Graminoid | Perennial |
| POACEAE | <i>Eragrostis trichophora</i> Coss. & Durieu | LC | No | Graminoid | Perennial |
| POACEAE | <i>Fingerhuthia africana</i> Lehm. | LC | No | Graminoid | Perennial (occ. annual) |
| POACEAE | <i>Heteropogon contortus</i> (L.) Roem. & Schult. | LC | No | Graminoid | Perennial |
| POACEAE | * <i>Lamarckia aurea</i> (L.) Moench | Not Evaluated | No | Graminoid | Annual |
| POACEAE | <i>Leptochloa fusca</i> (L.) Kunth | LC | No | Graminoid | Perennial |
| POACEAE | <i>Melinis repens</i> (Willd.) Zizka subsp. <i>repens</i> | LC | No | Graminoid | Annual (occ. perennial) |
| POACEAE | <i>Panicum coloratum</i> L. var. <i>coloratum</i> | LC | No | Graminoid | Perennial |
| POACEAE | <i>Panicum maximum</i> Jacq. | LC | No | Graminoid | Perennial |
| POACEAE | <i>Pogonarthria squarrosa</i> (Roem. & Schult.) Pilg. | LC | No | Graminoid | Perennial (occ. annual) |
| POACEAE | <i>Schmidtia pappophoroides</i> Steud. | LC | No | Graminoid | Perennial |
| POACEAE | <i>Setaria verticillata</i> (L.) P.Beauv. | LC | No | Graminoid | Annual |

| Family | Species | Threat status | SA Endemic | Growth forms | Lifecycle |
|---------------|--|---------------|------------|------------------------------|-------------------------|
| POACEAE | <i>Sporobolus fimbriatus</i> (Trin.) Nees | LC | No | Graminoid | Perennial |
| POACEAE | <i>Stipagrostis uniplumis</i> (Licht.) De Winter var. <i>uniplumis</i> | LC | No | Graminoid | Perennial (occ. annual) |
| POACEAE | <i>Tragus berteronianus</i> Schult. | LC | No | Graminoid | Annual |
| POACEAE | <i>Tragus racemosus</i> (L.) All. | LC | No | Graminoid | Annual |
| POACEAE | <i>Trichoneura grandiglumis</i> (Nees) Ekman | LC | No | Graminoid | Perennial |
| POACEAE | <i>Urochloa panicoides</i> P.Beauv. | | No | Graminoid | Annual |
| POACEAE | <i>Urochloa stolonifera</i> (Gooss.) Chippind. | LC | No | Graminoid | Perennial |
| POACEAE | <i>Anthephora pubescens</i> Nees | LC | No | Graminoid | Perennial |
| POACEAE | <i>Brachiaria marlothii</i> (Hack.) Stent | LC | No | Graminoid | Annual (occ. perennial) |
| POACEAE | <i>Coelachyrum yemenicum</i> (Schweinf.) S.M.Phillips | LC | No | Graminoid | Perennial |
| POACEAE | <i>Digitaria eriantha</i> Steud. | LC | No | Graminoid | Perennial |
| POACEAE | <i>Enneapogon desvauxii</i> P.Beauv. | LC | No | Graminoid | Annual (occ. perennial) |
| POACEAE | <i>Eragrostis echinochloidea</i> Stapf | LC | No | Graminoid | Perennial |
| POACEAE | <i>Eustachys paspaloides</i> (Vahl) Lanza & Mattei | LC | No | Graminoid | Perennial |
| POACEAE | <i>Oropetium capense</i> Stapf | LC | No | Graminoid | Perennial |
| POACEAE | <i>Sporobolus acinifolius</i> Stapf | LC | No | Graminoid | Perennial |
| POACEAE | <i>Sporobolus ioclados</i> (Trin.) Nees | LC | No | Graminoid | Perennial |
| POACEAE | <i>Stipagrostis obtusa</i> (Delile) Nees | LC | No | Graminoid | Perennial |
| POACEAE | <i>Triraphis andropogonoides</i> (Steud.) E.Phillips | LC | No | Graminoid | Perennial |
| POLYGALACEAE | <i>Polygala leptophylla</i> Burch. var. <i>armata</i> (Chodat) Paiva | LC | No | Dwarf shrub | Perennial |
| PORTULACACEAE | <i>Portulaca hereroensis</i> Schinz | LC | No | Herb, succulent | Annual |
| PORTULACACEAE | <i>Portulaca kermesina</i> N.E.Br. | LC | No | Herb, succulent | Annual (occ. perennial) |
| RHAMNACEAE | <i>Helinus spartioides</i> (Engl.) Schinz ex Engl. | LC | No | Dwarf shrub | Perennial |
| SANTALACEAE | <i>Thesium hystricoides</i> A.W.Hill | LC | No | Dwarf shrub, parasite, shrub | Perennial |

| Family | Species | Threat status | SA Endemic | Growth forms | Lifecycle |
|------------------|--|---------------|------------|------------------------------|-------------------------|
| SANTALACEAE | <i>Thesium hystrix</i> A.W.Hill | LC | No | Dwarf shrub, parasite, shrub | Perennial |
| SCROPHULARIACEAE | <i>Aptosimum albomarginatum</i> Marloth & Engl. | LC | No | Dwarf shrub | Perennial |
| SCROPHULARIACEAE | <i>Jamesbrittenia atropurpurea</i> (Benth.) Hilliard subsp. <i>atropurpurea</i> | LC | No | Dwarf shrub, shrub | Perennial |
| SCROPHULARIACEAE | <i>Aptosimum elongatum</i> Engl. | LC | No | Dwarf shrub | Perennial |
| SCROPHULARIACEAE | <i>Aptosimum junceum</i> (Hiern) Philcox | LC | No | Dwarf shrub, shrub | Perennial |
| SCROPHULARIACEAE | <i>Aptosimum lineare</i> Marloth & Engl. var. <i>lineare</i> | LC | No | Dwarf shrub | Perennial |
| SCROPHULARIACEAE | <i>Peliostomum leucorrhizum</i> E.Mey. ex Benth. | LC | No | Dwarf shrub | Perennial |
| SOLANACEAE | <i>Solanum burchellii</i> Dunal | LC | No | Dwarf shrub, shrub | Perennial |
| SOLANACEAE | <i>Solanum lichtensteinii</i> Willd. | LC | No | Dwarf shrub, shrub | Perennial |
| SOLANACEAE | <i>Solanum tomentosum</i> L. var. <i>tomentosum</i> | LC | No | Dwarf shrub, herb, shrub | Perennial |
| SOLANACEAE | * <i>Datura stramonium</i> L. | Not Evaluated | No | Herb, shrub | Annual |
| SOLANACEAE | <i>Lycium cinereum</i> Thunb. | LC | No | Dwarf shrub, shrub | Perennial |
| SOLANACEAE | <i>Lycium hirsutum</i> Dunal | LC | No | Dwarf shrub, shrub | Perennial |
| SOLANACEAE | <i>Lycium pilifolium</i> C.H.Wright | LC | No | Dwarf shrub, shrub | Perennial |
| SOLANACEAE | <i>Solanum catombelense</i> Peyr. | LC | No | Dwarf shrub, shrub | Perennial |
| VAHLIACEAE | <i>Vahlia capensis</i> (L.f.) Thunb. subsp. <i>vulgaris</i> Bridson var. <i>linearis</i> E.Mey. ex Bridson | LC | No | Herb | Annual (occ. perennial) |
| VERBENACEAE | <i>Chascanum hederaceum</i> (Sond.) Moldenke var. <i>hederaceum</i> | LC | No | Herb | Perennial |
| VERBENACEAE | <i>Chascanum pinnatifidum</i> (L.f.) E.Mey. var. <i>pinnatifidum</i> | LC | No | Herb | Perennial |
| VERBENACEAE | <i>Lantana rugosa</i> Thunb. | LC | No | Shrub | Perennial |
| ZYGOPHYLLACEAE | <i>Tribulus excrucians</i> Wawra | | No | Dwarf shrub, shrub | Biennial |
| ZYGOPHYLLACEAE | <i>Tribulus terrestris</i> L. | LC | No | Herb | Annual |
| ZYGOPHYLLACEAE | <i>Tribulus zeyheri</i> Sond. subsp. <i>zeyheri</i> | LC | No | Dwarf shrub, herb | Perennial |

13 APPENDIX C: Recorded positions of red data or protected species

Table 13-1: Coordinates of some recorded ToPS, red data and protected plant species

Note: The numbered labels on Figure 11 (p. 27) correspond to the serial number (S/N) in the first column of Table 13-1.

| S/N | SPECIES | Coordinates | | No of Specimens |
|----------------|--|----------------|----------------|-----------------|
| | | S | E | |
| Preferred Site | | | | |
| 1 | <i>Harpagophytum procumbens</i> subsp. <i>procumbens</i> | 27° 15' 56.0"S | 23° 01' 05.1"E | 5 |
| 2 | <i>Ruschia</i> species | 27° 14' 53.2"S | 23° 01' 36.1"E | ±20 |
| 3 | <i>Ruschia</i> species | 27° 14' 53.1"S | 23° 01' 32.0"E | |
| 4 | <i>Ruschia</i> species | 27° 15' 01.5"S | 23° 01' 36.8"E | 5 |
| | <i>Harpagophytum procumbens</i> subsp. <i>procumbens</i> | | | 5 |
| 5 | <i>Harpagophytum procumbens</i> subsp. <i>procumbens</i> | 27° 15' 04.7"S | 23° 01' 37.3"E | 4 |
| 6 | <i>Harpagophytum procumbens</i> subsp. <i>procumbens</i> | 27° 15' 27.9"S | 23° 01' 28.0"E | 3 |
| 7 | <i>Harpagophytum procumbens</i> subsp. <i>procumbens</i> | 27° 15' 51.2"S | 23° 01' 02.8"E | ±10 |
| 8 | <i>Harpagophytum procumbens</i> subsp. <i>procumbens</i> | 27° 15' 57.6"S | 23° 00' 51.9"E | ±10 |
| 9 | <i>Harpagophytum procumbens</i> subsp. <i>procumbens</i> | 27° 15' 35.8"S | 23° 00' 45.0"E | 3 |
| 10 | <i>Harpagophytum procumbens</i> subsp. <i>procumbens</i> | 27° 14' 36.3"S | 23° 01' 25.4"E | ±40 |
| | <i>Boophone disticha</i> | 27° 14' 36.6"S | 23° 01' 25.8"E | 1 |
| 11 | <i>Harpagophytum procumbens</i> subsp. <i>procumbens</i> | 27° 14' 33.2"S | 23° 01' 23.1"E | ±60 |
| 12 | <i>Nerine laticoma</i> | 27° 15' 13.9"S | 23° 01' 52.6"E | ±5 |
| 13 | <i>Nerine laticoma</i> | 27° 15' 16.0"S | 23° 01' 50.8"E | ±30 |
| 14 | <i>Euphorbia c.f. braunsii</i> | 27° 15' 38.3"S | 23° 01' 33.1"E | 1 |

Table 13-2: Calculations of protected tree density in the study area

| Veg Unit | Transect No. | Species frequency (as counted on 4000 m ²) | | | | | |
|-----------------------------------|---------------|--|--------------|--------------|----------------------------|-------------|--------------|
| | | <i>Acacia erioloba</i> | | | <i>Acacia haematoxylon</i> | | |
| | | 1 to <2 m | >2 m | Total | 1 to <2 m | >2 m | Total |
| 1 (183ha) | 1 | 6 | 1 | 7 | 23 | 3 | 26 |
| | 2 | 3 | 3 | 6 | 41 | 3 | 44 |
| | 3 | 6 | 2 | 8 | 31 | 1 | 32 |
| | 4 | 1 | 3 | 4 | 10 | 0 | 10 |
| | 5 | 0 | 2 | 2 | 62 | 4 | 66 |
| | 6 | 6 | 5 | 11 | 8 | 3 | 11 |
| | 7 | 1 | 1 | 2 | 17 | 1 | 18 |
| | Ave: | 3.3 | 2.4 | 5.7 | 27.4 | 2.1 | 29.6 |
| 2 (122ha) | 1 | 0 | 15 | 15 | 0 | 0 | 0 |
| | 2 | 2 | 12 | 14 | 0 | 1 | 1 |
| | 3 | 2 | 16 | 18 | 0 | 0 | 0 |
| | 4 | 6 | 18 | 24 | 0 | 0 | 0 |
| | 5 | 5 | 19 | 24 | 6 | 3 | 9 |
| | 6 | 5 | 19 | 24 | 5 | 0 | 5 |
| | 7 | 5 | 31 | 36 | 2 | 2 | 4 |
| | Ave: | 3.6 | 18.6 | 22.1 | 1.9 | 0.9 | 2.7 |
| Species density / ha | | | | | | | |
| 1 (183ha) | 1 | 15 | 2.5 | 17.5 | 57.5 | 7.5 | 65 |
| | 2 | 7.5 | 7.5 | 15 | 102.5 | 7.5 | 110 |
| | 3 | 15 | 5 | 20 | 77.5 | 2.5 | 80 |
| | 4 | 2.5 | 7.5 | 10 | 25 | 0 | 25 |
| | 5 | 0 | 5 | 5 | 155 | 10 | 165 |
| | 6 | 15 | 12.5 | 27.5 | 20 | 7.5 | 27.5 |
| | 7 | 2.5 | 2.5 | 5 | 42.5 | 2.5 | 45 |
| | Ave: | 8.21 | 6.07 | 14.29 | 68.57 | 5.36 | 73.93 |
| 2 (122ha) | 1 | 0 | 37.5 | 37.5 | 0 | 0 | 0 |
| | 2 | 5 | 30 | 35 | 0 | 2.5 | 2.5 |
| | 3 | 5 | 40 | 45 | 0 | 0 | 0 |
| | 4 | 15 | 45 | 60 | 0 | 0 | 0 |
| | 5 | 12.5 | 47.5 | 60 | 15 | 7.5 | 22.5 |
| | 6 | 12.5 | 47.5 | 60 | 12.5 | 0 | 12.5 |
| | 7 | 12.5 | 77.5 | 90 | 5 | 5 | 10 |
| | Ave: | 8.93 | 46.43 | 55.36 | 4.64 | 2.14 | 6.79 |
| Number of specimens per VU | | | | | | | |
| 1 | | 1503 | 1111 | 2614 | 12549 | 980 | 13529 |
| 2 | | 1089 | 5664 | 6754 | 566 | 261 | 828 |
| | Total: | | | 9368 | | | 14357 |