

Effects of phytohormones and alternative sources on the propagation of *Ternstroemia cameroonensis* Cheek. by marcotting in the Lebialem Highlands, Cameroon.

factors influencing *Ternstroemia cameroonensis* Cheek. propagation in Cameroon's Lebialem Highlands

ABSTRACT

Marcotting is a method of vegetative propagation which still finds its relevance in the present day forest management and routine practice to produce viable seedlings for the regeneration. Of *T. cameroonensis*. There is dearth of information regarding the propagation of *T. cameroonensis* by marcotting. Hence effects of phytohormone application and alternative sources on the propagation of *T. cameroonensis* by marcotting were investigated in the Lebialem Highlands. Marcotts had an overall survival percentage of 35.41% (conditions culture??). you should give a brief overview of the tests carried out and the parameters to be studied Those established at the middle of the crown had the highest survival percentage (15.27%) followed by those at the lower (11.80 %). Also marcotts established on branches with larger diameter (4-6 cm) and shorter length (50 cm) had the highest survival percentage (20.13 and 19.44 respectively). Again cuttings treated with IBA (0.5 g/l and 0.2 g/l) and CW (4 h) had the best performance (for what??) compared to other pre-treatments. According to the obtained results, ~~From our findings,~~ *T. cameroonensis* can be amenable through marcotts.

Formatted: Highlight

Formatted: Strikethrough

Keywords: *Ternstroemia cameroonensis*, phytohormone and alternative sources, rooting of marcotts, branch diameter and length, marcottst position.

1. INTRODUCTION

Ternstroemia cameroonensis is a ~~critically~~ highly endangered medicinal plant species found only ~~endemic~~ in the Lebialem highlands [1; 2]. It growing 15-18m tall with inflorescence of four to five single flowers on the end of a stem, oval fruits and leathery leaves 6-7 x 22.5 [3].

Formatted: Strikethrough

Formatted: Strikethrough

This plant is of pharmaceutical importance to the indigenous people as decoction of bark is used to treat anaemia, stomach ache, vomiting, urinary tract infections, infertility and epilepsy [4]. A poster depicting the species, for use in Cameroon and promoting its conservation was produced and distributed by Kew [5]. Tacham *et al.* [6], report that the species is overexploited for sale locally. No regeneration has been detected possibly because any seedlings that develop might be swept away by run-off down the steep slopes on which the surviving trees grow [2]. The work of Nkemnkeng *et al.* [7] indicate that the species is highly vulnerable in the Lebialem highland due method of exploitation with few mature individuals in the wild. The species has not been cultivated nor seed-banked. Seeds of some species of *Ternstroemia* have been shown to be dormant, particularly *Ternstroemia washikiatii* germinating after six to seven months [8]. Recent study of Nkemnkeng *et al.* [9] indicates that propagation of *T. cameroonensis* through seed and stem cuttings were inappropriate. However, a great deal of work is required to fully develop the propagation potential of this species.

Asexual propagation describes the regeneration and multiplication of plant from vegetative parts, like buds, leaves, single cells/tissues as well as cuttings of roots and stems [10]. It offers a wide array of applications in indigenous fruit tree domestication efforts and general plant genetic resources conservation programmes. With these applications, experts are able to select desired characters available in the wild tree population by fixating the genetic variation of such trees in the natural stands. Ultimately, the objective is to multiply large quantity of improved seedlings for interested farmers and other users [11, 12]. Marcotting is a vegetative propagation technique characterized by the initiation of adventitious roots on one part of the tree branch in situ. After root initiation, the marcotts is weaned from the tree and transplanted in a substrate where it grows independently of the mother tree. Like other vegetative propagation techniques, the main advantages of marcotting are cloning selected trees with desirable traits and shortening the period for fruit production [13, 14].

Diverse factors (genetic, environmental and physiological) may influence the rooting and shooting abilities of marcotts. Studies had shown that rooting ability of marcotts in some species may vary with marcott length, branch diameter, bark thickness, rooting substrate, hormone type and concentration [13, 15]. It has also been demonstrated that rooting and sprouting requirements are highly variable within and among species. Therefore, assessing factors that affect rooting and sprouting abilities of marcotts in each species is an important prerequisite for its successful propagation through air-layering. This [purpose of this](#) study

was ~~to determine~~, meant to assess the effect of phytohormones as well as alternative sources, branch diameter, branch length and the marcott's position in the crown on rooting and survival of weaned marcotts.

Formatted: Strikethrough

2. METHODOLOGY

2.1. Study site

This study was carried out in two phases. The first phase in the Lebialem Highlands located in the North Eastern part of the Southwest Region of Cameroon between latitude 5°38 N and 5°43 N and longitude 9°58 E and 10°27 E [16].

The climate of this region is that of the Cameroon mountain range characterized by high winds and low sunshine. The average daily temperature varies very much with season and ranges from 17°C to 32°C [17]. The area is between altitudes 1000 m to 2700 m above sea level and annual rainfall is 2222 mm.

2.2. Marcotts preparation and setting

Trees were selected in the wild considering areas which are less prone to anthropogenic activities. On each selected tree marcotts were set at each level that is lower, middle and upper section of the crown. These levels consisted the blocks within which were treatment and repetitions on different tree. The marcotts were set up is a 2x2x4 factorial design with 3 levels and 3 repetitions on different trees that is a total of 144 marcotts were setup for this trial. Marcotts were set up in August on orthotropic and oblique oriented branches in the morning ~~why???~~. The bark was completely stripped off with a knife 5 cm away from the crotch (formed by two branches) and over a length of 5 to 10 cm. After application of hormone and coconut water on the stripped area, it was covered with moist decomposed sawdust wrapped in a transparent plastic sheet and secured at each end with a rubber band [14]. Three experiments were setup with details presented below (Figure 1).

- ❖ Experiment 1: Effect of IAA, IBA and Coconut water on marcotts' rooting abilities

~~In this experyience, t~~Three trees were used ~~for this experiment~~. On each tree marcotts setting were ~~taken~~ at random in the crown. After stripping, rooting reagents were applied and ~~the plant were~~ covered with moist decomposed sawdust. A total of 48 marcotts were ~~planted, set~~ per tree, ~~with IAA and IAA~~ applied to 12 ~~marcotts of them~~, IBA ~~to another on~~ 12, CW ~~also on~~ applied to 12, and 12 marcotts ~~left~~ and 12 marcotts were untreated (control).

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

- ❖ Experiment 2: Effect of marcotts's position on rooting ability

Three trees were chosen and on each, the crown was divided into the lower, middle and upper sections. In each section, four marcotts were set that is untreated (control), IAA, IBA and CW. [to be rephrased](#)

Formatted: Highlight

❖ Experiment 3: The effect of branch length and diameter on marcotts' survival

Marcotts were established on branches with different branch length and diameter. Branch length and diameter were within different ranges that is 40-60 cm (L₅₀), 80-120 cm (L₁₀₀) and 2-4 cm (D₃), 4-6 cm (D₆) respectively [on what basis you chose the lengths and diameters of the branches???](#). Equally after stripping, rooting reagent were applied and covered with decomposed sawdust using polythene paper and tied at both ends. The marcottsts were watered with about 20 ml of water every 2 weeks using a 10 ml syringe. The experimental design was randomized complete block. Assessment of dependent variables that is rooted marcotts, unrooted marcotts and mortality were evaluated every after 14 weeks for a period of 10 months.

2.3. Data collection.

This consisted of mortality rate, number of rooted marcotts as well as root length. All these were evaluated every after 12 weeks for a period of 10 months.

2.4. Data analysis.

Data collected were entered into excel and subjected into various analysis.

Percentage of rooting of marcotts as well as number of roots, rate of mortality and survival percentages were analysed. Early growth parameters of marcotts were subjected to Analysis of Variance (ANOVA) using the statistical programme XLStat, 2016 where the least significant differences (LSD) between the mean were detected and separated at $p \leq 0.05$.

3. RESULT

3.1 Effects of coconut water (CW) and hormones (IAA and IBA) marcotts rooting of *T. cameroonensis*.

Out of the 144 marcotts established, 51 survived after a period of 8 months given a survival percentage of 35.41%. Those established at the middle of the crown had the highest survival percentage (15.27%) followed by those at the base (11.80 %) and lastly by those at the top (8.33%). Also marcotts established on branches with larger diameter (4-6 cm) and shorter length (50cm) had the highest survival percentage that is 20.13% and 19.44 respectively while those with smaller diameter and longer branch length had the least. Again cuttings treated with IBA and CW had the best performance in terms of survival, number of roots and root length compared to other treatments. [you are talking more about percentage of survival](#)

than rooting, you should highlight the marcotts having regenerated roots and their percentages???

3.2 Effect of coconut water, and hormones on rooting ability, and marcotts performance

There was a significant difference in the survival percentage, number of roots and roots length of marcotts with respect to pre-treatment. IBA had the best performance (47.22%, 2.33 and 2.76 respectively) in all the parameters followed by CW (41.57%, 0.83 and 1.33) while the control treatment was the least. The percentages are not clear. How did you calculate the percentages of roots marcotts? based on each number (initial number of marcotts or just survival ones)? What about the length of the roots? This paragraph should be revised.-(table 1).

Formatted: Highlight

3.3 Effect of branch diameter and branch length on rooting ability and marcotts performance

Branch length and diameter had no significant effect on the survival percentages of marcotts. This is also observed in branch length and diameter of 80-120 cm and 2-3 cm respectively explain???. However, branch length and diameter within 40 – 60 cm and 4-6 cm significantly affected rooting of marcotts. Marcotts of branch length within 40-60 cm had the highest number of leaves ???? and root length (1.44 cm and 1.64 cm respectively) while those of branch length within 80-120 cm had the least number of roots and root length (0.167, 0.083 respectively) (table 2). In addition, marcotts of branch diameter within 4-6 cm had the highest number of leaves ??? and root length (1.72 cm and 1.972 cm respectively) while those of branch length within 2-3 cm had the least number of roots and root length (0.056) (Table 3).

3.4 Effect of marcotts position on rooting ability and marcotts performance

Marcotts position did not significantly affected survival percentage, as well as number of roots and root length explain?. Despite this, marcotts established at middle of the tree crown had better performance in terms of survival percentages, number of root and root length % than those in lower and upper positions % give -examples (table 4, 5 and 6).

3.5 Survival of marcotts of *T. cameroonensis*

Rooted marcotts of *T. cameroonensis* were weaned into polythene bags at the nursery (Figure 26). Out of 51 rooted marcotts weaned, only seven survived giving a percentage of 13.72% after a period of two months.

4. DISCUSSION

Ascertaining the suitability of marcotting or any other vegetative propagation technique for that matter for the propagation of any particular species before recommendation for large

scale use is very important to avoid the loss of tree stands to mortality due to incompatibility. The results are however in agreement with the findings of Steward, [18] who reported that variation in the rooting of marcotts is a factor that can be attributed to environmental conditions that increases the chances of marcotts or mother tree being susceptible to varying degrees of stress and therefore different levels of response to the marcotting procedure. Marcotts within the diameter class of 4-6 cm performed better maybe due to high carbohydrate and auxin reserves. This ~~corroborate~~corroborates with the results of Assah, [19]; Okonkwo *et al.* [20] on vegetative propagation of *Dacryodes edulis*, *Garcinia kola* respectively through marcotting. This result is supported by the work of Menzel, [21] who reported variations in root growth in marcotted trees of *Artocarpus altilis*, and the findings of Kengue, [14] and Mialoundama *et al.*, [15] who found differences in root volume in marcotted trees of *Dacryodes edulis* and *Ricinodendron heudelotii* respectively and associated the variations to differences in size of marcotted branches. This is in agreement with Brian and Nina, [22] who postulated that large diameter tree branches tend to have higher carbohydrate and auxin reserves than smaller ones and are therefore more likely to root better than their smaller counterparts. The benefit of the present study is that it reveals that the high carbohydrate reserves in larger diameter branches that promotes rooting also promotes and supports the survival of marcotts while still attached to the mother plant.

In addition, marcotts position equally had an influence on rooting ability, number of roots and root length. Those at the middle of the crown rooted better maybe due the high humid content compared to lower part as well as upper parts of the crown. This is apparent with the findings of Ogbu *et al.* [23] that adventitious root formation by marcotted branches of *P. macrophylla* required high humid condition which is naturally guaranteed during the peak of wet season weather, and vice versa. Obviously, this behaviour may be attributed to the prevailing high moisture and cool temperature associated with the period of peak rainfall. It has been long established that root formation is influenced by auxins produced by the plant itself [10; 24; 25]. The physiology of tree growth phases show that as plant undergoes active vegetative growth cycle; buds release endogenous auxin that moves basipetally to stimulate root production [24; 25]. So by implication and in response to the deliberate wound caused by girdling in marcotting operation, branches of the species began formation of adventitious roots. This is an affirmation of similar reports by Awodoyin and Olaniyan [24], Leakey [26] and Hartmann *et al.* [10] that infliction of wounds on stem brings about auxin induced stimulation of vascular meristematic tissues to form adventitious roots in cuttings and layering techniques.

5. CONCLUSION.

Rooting ability of *T. cameroonensis* marcotts proved successful in phytohormones (IAA and IBA) as well as alternative sources (coconut water). IBA had the best performance followed by coconut water while IAA was the least. Thus, rooting of marcotts of *Ternstroemia cameroonensis* with plant growth regulator (IBA, CW) presents a viable propagation technique to be used in enrichment planting programmed of this important critically endangered medicinal plant, though weaned marcotts had a very low survival percentage.

outlook:

improve the success rate to ensure a high percentage of survival rooted marcotts!!!!

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES.

1. IUCN. IUCN Red List categories and criteria: Version 3.1. Second edition. Gland, Switzerland and Cambridge, UK: IUCN. 2012; Available from: <http://www.iucnredlist.org/>, [accessed: Jan. 2020].
2. Cheek M, Tchiengue B, Tacham WN. *Ternstroemia cameroonensis* (Ternstroemiaceae), a new medicinally important species of montane tree, nearly extinct in the Highlands of Cameroon. *Blumea*. 2017; **62**:53–57.
3. WHINCONET. Plant and Animal Guide for the Western Cameroon Highlands, 14p. Tropical trees: the potential for domestication and the rebuilding of forest resources. London: HMSO. 2007; 15p.
4. Focho DA, Ndam WT, Fonge BA. Medicinal plants of Aguambu—Bamumbu in the Lebalem highlands, southwest province of Cameroon. *African Journal of Pharmacy and Pharmacology*. 2009; **3**(1): 1–13.
5. Cheek M. *Ternstroemia polypetala* poster. Royal Bontanic Gardens, Kew. *Ternstroemia* (*Pentaphylacaceae*) from La Amistad Binational Park and World Heritage Property, Costa Rica and Panama. *Phytotaxa*. 2000; **217**: 87–91.

Formatted: Highlight

Formatted: Font: (Default) +Body, 11 pt, No underline, Underline color: Auto, Font color: Auto, Highlight

Formatted: Highlight

6. Tacham WN, Fonge BN, Fonkou T. Traditional medicine and ethnobotanical use of wild plants by the Mundani people of Wabane, South West Region, Cameroon. *Journal of Ethnobiology and Traditional Medicine. Photon.* 2015; **125**:1060–1080.
7. Nkemnkeng FG, Anjah GM, Tacham WN, Mbogue CNN, Wouokoue JBT, Nguetsop VF. Population distribution, conservation status and vulnerability of *Ternstroemia cameroonensis* Cheek. in the Lebialem Highlands, Cameroon. *International Journal of Current Research in Bioscience and Plant Biology.* 2021; **8**;(2), 1-10.
8. Xavier C, Carmen UU. *Ternstroemia washikiatii* (Pentaphylacaceae): a new tree species from Eastern Ecuador. *Harvard Papers in Botany.* 2016; 21(2):337–339.
9. Nkemnkeng FJ, Anjah GM, Tacham WN, Mbogue CNN, Nguetsop VF. Propagation of *Ternstroemia cameroonensis*: an approach towards the conservation of a critically endangered medicinal plant species in the Lebialem Highlands, Cameroon. *Journal of Biodiversity and Environmental Sciences.* 2021; (In Review).
10. Hartmann HT, Kester DE, Davies Jr. FT, Geneve RL. Plant Propagation: principles and practices (7th ed.). Prentice- Hall Inc., New Delhi. 2007; 293 – 603 p.
11. Tchoundjeu Z, De-Wolf J, Jaenicke H. Vegetative propagation for domestication of agroforestry trees. *Agroforestry Today.* 1997; 9(2): 10-12.
12. Awodoyin RO, Olubode OS, Ogbu JU, Balogun RB, Nwawuisi JU, Orji KO. Indigenous Fruit Trees of Tropical Africa: Status, Opportunity for Development and Biodiversity Management. *Agricultural Sciences.* 2015; 6: 31-41. <http://dx.doi.org/10.4236/as.2015.61004>.
13. Hartmann HT, Kester DE, Davies FT, Geneve RL. Plant propagation: Principle and Practices. Sixth Edition. Prentice Hall. 1997; 239-513 p.
14. Kengue J. Safou, *Dacryodes edulis* (G. Dom) H. J. Lam. ICUC, RPM Reprographics, Chicghester, Angleterre. 2002; 144 p.
15. Mialoundama F, Avana ML, Youmbi-EMampouya PC, Tchoundjeu Z, Mbeuyo M, Galamo GR, Bell JM, Kogpuep F, Tsobeng AC, Abega J. Vegetative propagation of *Dacryodes edulis* (G. Don) H. Lam by marcotts, cuttings and micro-propagation. *Journal of Forests Trees and livelihoods.* 2002; 12(1):85-96.
16. Nkembi, L. Comparative study of community and government patrols in enhancing sustainable wildlife conservation in the Banyang-Mbo sanctuary, Cameroon. Report for the Ministry of Environment and Forestry. *MINEF.* 2004; Unpublished, 14 p.
17. Zogning, C.N., Tiafack, O. The catastrophic geomorphological processes in humid tropical Africa: a case study of the recent land slide disasters in Cameroon. *Sedimentary Geology.* 2007; 199(1-2):13–27.

Formatted: Default Paragraph Font, Font: (Default) +Body, 11 pt, Highlight

Formatted: No underline, Font color: Auto, Highlight

Formatted: Highlight

Formatted: No underline, Font color: Auto, Highlight

Formatted: Highlight

Formatted: No underline, Font color: Auto, Highlight

Formatted: Highlight

Formatted: No underline, Font color: Auto, Highlight

Formatted: Highlight

Formatted: No underline, Font color: Auto, Highlight

Formatted: Highlight

Formatted: No underline, Font color: Auto, Highlight

Formatted: Highlight

18. Steward A. Let's propagate: A plant propagation manual for Australia. ABC books Sydney, Australia. 2012; 240 p.

Formatted: No underline, Font color: Auto, Highlight

Formatted: Highlight

19. Asaah EK. Beyond vegetative propagation of indigenous fruit trees: case study of *Dacryodes edulis* (G. Don) H. J. Lam and *Allanblackia floribunda* Oliv. PhD. Thesis, Faculty of Bioscience Engineering Ghent University, Belgium. 2012; 231 p.

Formatted: No underline, Font color: Auto, Highlight

Formatted: Highlight

20. Menzel, C. M. 1991. Litchi chinensis Sonn. In: Verheij, E. W. M, and Coronel, R. E., (eds.), Plant Resources of South-East Asia. Pudoe, Wageningen, the Netherlands: 191-195.

Formatted: No underline, Font color: Auto, Highlight

Formatted: Highlight

21. Okonkwo HO, Koyejo OA, Oyediran RI (2019) Pre-Severance and Post-Severance Survival, Growth and Fruiting of *Allanblackia floribunda* Oliv. Marcotts. *World Journal of Agricultural Sciences* 15 (4): 215-219.

Formatted: No underline, Font color: Auto, Highlight

Formatted: Highlight

22. Brian KM, Nina LB. Etiolation and banding effects on adventitious root formation. In Davis T.D., Haissig BE., Sankhla N. (eds): Adventitious root formation in cuttings. Advances in Plant Sciences Series, 2. Dioscorides press. 1988; 30-43 p.

Formatted: No underline, Font color: Auto, Highlight

Formatted: Highlight

23. Ogbu JU, Awodoyin RO, Isienyi NC. Effect of phytohormone and phenology on domestication of *Pentaclethra macrophylla* benth. by marcotting in derived savanna zone of southeast nigeria. *Nigerian agricultural journal*. 2019; 50(2):18-23.

Formatted: No underline, Font color: Auto, Highlight

Formatted: Highlight

24. Awodoyin RO, Olaniyan AA. Air layering in clonal propagation of Guava (*Psidium guajava* L.): Effects of season and IBA growth hormone on root production. Proceedings of 18th Annual Conference of the Horticultural Society of Nigeria. 2000; 113 – 116.

Formatted: No underline, Font color: Auto, Highlight

Formatted: Highlight

25. Jaenicke H, Beniast J. Vegetative tree propagation in agroforestry. ICRAF, Nairobi, Kenya. 2002; 1 – 30., 75 – 82.

Formatted: No underline, Font color: Auto, Highlight

Formatted: Highlight

26. Leakey RRB. Physiology of vegetative reproduction. In: Burley J, Evans E, Younquist JA, editors. Encyclopaedia of forest sciences. London: Academic Press. 2004; 1655–1668 p.

Formatted: No underline, Font color: Auto, Highlight

Tables

Table 1: Effect of coconut water and hormones on rooting ability and marcots performance.

Treatment	Sur%	NR	RL
Control	27.78 ^a	0.389 ^a	0.305 ^a
CW	41.67 ^{ab}	0.83 ^a	1.33 ^{ab}
IBA	47.22 ^b	2.33 ^b	2.76 ^b
IAA	25.0 ^a	0.5 ^a	0.56 ^a

Sur%: survival percentage, NR: number of roots, RL: root length

[Statistic ????: a,b,c,d,e Means +- SE with different letters in the same column are **high significance P<0.001 significant *p<0.05](#)

Table 2: Effect of branch length on rooting ability and marcots performance

L	SUR%50	SUR%100	NR 50	NR100	RL.50	RL 100
Control	33.33 ^a	22.22 ^a	0.22 ^a	0.167 ^a	0.22 ^a	0.083 ^a
CW	44.44 ^a	38.89 ^a	0.389 ^{ab}	0.44 ^a	0.44 ^{ab}	0.89 ^a
IBA	50.0 ^a	44.44 ^a	1.44 ^b	0.89 ^a	1.64 ^b	1.125 ^a
IAA	27.78 ^a	22.22 ^a	0.278 ^a	0.22 ^a	0.22 ^a	0.33 ^a

50, 100: range of branch length (40-60 cm and 80-120 cm respectively)

Sur%: survival percentage, NR: number of roots, RL: root length

[Statistic ????: a,b,c,d,e Means +- SE with different letters in the same column are **high significance P<0.001 significant *p<0.05](#)

Table 3: Effect of branch diameter on rooting ability and marcotts performance

Formatted Table

Formatted: Space After: 0 pt, Line spacing: single

Formatted: Font: 11 pt, Bold

Formatted: Normal

Diameter	SUR.3	Sur.6	NR.3	NR.6	RL.3	RL.6
Control	22.22 ^a	33.33 ^a	0.056 ^a	0.33 ^a	0.11 ^a	0.194 ^a
CW	33.33 ^a	50.0 ^a	0.167 ^a	0.67 ^{ab}	0.25 ^a	1.083 ^{ab}
IBA	44.44 ^a	50.0 ^a	0.61 ^a	1.72 ^b	0.79 ^a	1.972 ^b
IAA	22.22 ^a	27.78 ^a	0.056 ^a	0.44 ^a	0.056 ^a	0.5 ^{ab}

3, 6: range of branch diameter (2-4 cm and 4-6 cm respectively)

Sur%: survival percentage, NR: number of roots, RL: root length

Statistic ???: a,b,c,d,e Means +- SE with different letters in the same column are **high significance $P \leq 0.001$ significant * $p \leq 0.05$

Formatted: Space After: 0 pt, Line spacing: single

Formatted Table

Formatted: Space After: 0 pt, Line spacing: single

Formatted: Space After: 0 pt, Line spacing: single

Formatted: Space After: 0 pt, Line spacing: single

Formatted: Space After: 0 pt, Line spacing: single

Formatted: Space Before: 0 pt, After: 0 pt

Formatted: Space After: 0 pt

Table 4: Effects of position on survival percentage of marcott

Treatment	D.6, L.50			D.3, L.50			D.6, L.100			D.3, L.100		
	L	M	U	L	M	U	L	M	U	L	M	U
Control	33.33 ^a	33.33 ^a	33.33 ^a	33.33 ^a	33.33 ^a	33.33 ^a	50.0 ^a	33.33 ^a	33.33 ^a	0 ^a	33.33 ^a	0 ^a
CW	66.67 ^a	33.33 ^a	33.33 ^a	33.33 ^a	33.33 ^a	33.33 ^a	33.33 ^a	66.67 ^a	33.33 ^a	33.33 ^a	33.33 ^a	33.33 ^a
IBA	66.67 ^a	66.67 ^a	33.33 ^a	33.33 ^a	66.67 ^a	33.33 ^a	33.33 ^a	66.67 ^a	33.33 ^a	33.33 ^a	66.67 ^a	33.33 ^a
IAA	33.33 ^a	33.33 ^a	33.33 ^a	33.33 ^a	33.33 ^a	0 ^a	33.33 ^a	33.33 ^a	0 ^a	33.33 ^a	33.33 ^a	0 ^a

L: Lower position, M: Middle position, U: Upper position.

Statistic ???: a,b,c,d,e Means +- SE with different letters in the same column are **high significance $P \leq 0.001$ significant * $p \leq 0.05$

Formatted: Space After: 0 pt

Table 5: Effects of position on number of leaves of rooted marcotts.

Treatment	D.6, L.50			D.3, L.50			D.6, L.100			D.3, L.100		
	L	M	U	L	M	U	L	M	U	L	M	U
Control	0.33 ^a	0.67 ^a	0 ^a	0 ^a	0.33 ^a	0 ^a	0.67 ^a	0 ^a	0.33 ^a	0 ^a	0 ^a	0 ^a
CW	0.67 ^a	0.67 ^a	0 ^a	0.33 ^a	0.67 ^a	0 ^a	1.0 ^a	1.0 ^a	0.67 ^a	0 ^a	0 ^a	0 ^a
IBA	0.67 ^a	4.67 ^a	0.67 ^a	0.67 ^a	1.67 ^a	0.33 ^a	1.33 ^a	2.67 ^a	0.33 ^a	0.67 ^a	0.33 ^a	0 ^a
IAA	0.33 ^a	1.0 ^a	0 ^a	0.33 ^a	0 ^a	0 ^a	0 ^a	1.33 ^a	0 ^a	0 ^a	0 ^a	0 ^a

L: Lower position, M: Middle position, U: Upper position

Statistic ???: a,b,c,d,e Means +- SE with different letters in the same column are **high significance $P \leq 0.001$ significant * $p \leq 0.05$

Formatted: No underline, Font color: Accent 1, English (U.S.)

Formatted: Space After: 0 pt

Formatted: Font: 11 pt

Formatted: Normal

Table 6: Effects of position on root length of rooted marcotts.

Treatment	D.6, L.50			D.3, L.50			D.6, L.100			D.3, L.100		
	L	M	U	L	M	U	L	M	U	L	M	U

Control	0.167 ^a	0.5 ^a	0 ^a	0 ^a	0.67 ^a	0 ^a	0.33 ^a	0 ^a	0.167 ^a	0 ^a	0 ^a	0 ^a
CW	0.33 ^a	0.83 ^a	0 ^a	0.167 ^a	1.33 ^a	0 ^a	0.67 ^a	4.33 ^a	0.33 ^a	0 ^a	0 ^a	0 ^a
IBA	0.5 ^a	5.67 ^a	0.5 ^a	0.5 ^a	2.33 ^a	0.33 ^a	0.83 ^a	4.0 ^a	0.33 ^a	1.0 ^a	0.583 ^a	0 ^a
IAA	0.33 ^a	0.67 ^a	0 ^a	0.33 ^a	0 ^a	0 ^a	0 ^a	2.0 ^a	0 ^a	0 ^a	0 ^a	0 ^a

L: Lower position, M: Middle position, U: Upper position

Statistic???: a,b,c,d,e Means +- SE with different letters in the same column are **high significance $P \leq 0.001$ significant * $p \leq 0.05$

Formatted: Space Before: 0 pt, After: 0 pt

Formatted: Space After: 0 pt

Figures



Figure 1: Marcots establishment and monitoring in the field; A: stripped branch; B and C: stripped branch covered with decomposed sawdust and tied with polythene paper; D: irrigation of marcots in the dry season. [Pictures not so clear!!!](#)




Figure 2: Characteristics of root emergence from marcots of *T. cameroonensis*; [observation made after ??? number of days](#) A: most marcots treated IAA and at upper position, B and C: marcots treated with IBA, CW and at middle and lower position. [Pictures not so clear!!!](#)

Formatted: No underline, Font color: Accent 1, English (U.S.)



C



Figure 3: Transplant of weaned marcots of *T. cameroonensis*; **A:** rooted marcotts transplanted into polythene bags; B and C: sprouted transplanted marcotts.

UNDER PEER REVIEW